BTR Mega Test and Discussion

Dr. Zainab Vora

1. A 54-year-old woman comes to the clinic due to difficulty hearing for the past few weeks. During the neurologic examination, the physician assesses her hearing using a vibrating tuning fork. The handle of the tuning fork is placed on her left mastoid process until the sound is no longer audible. The tines are then quickly placed near the patient's left auditory meatus, and she reports hearing no sound. When the handle of the vibrating fork is placed on the middle of her forehead, she hears the vibration more strongly in her left ear than her right. This patient is most likely experiencing which of the following types of hearing loss?

A. Conductive loss in left ear
B. Conductive loss in right ear
C. Sensorineural loss in both ears
D. Sensorineural loss in left ear

2. Which of the following method of disinfection and sterilization can also kill spores?

1. Glutaraldehyde
2. Ethylene oxide
3. Pasteurization
4. Orthophthalaldehyde

A. 1, 3 and 4
B. 1, 2 and 4
C. 2,3 and 4
D. 1,2,3 and 4

Disinfectant	Activity	Advantages	Disadvantages	Recommendation
Glutaraldehyde	Broad spectrum microbicidal and sporicidal	Good compatibility	Requires activation Produces irritant fumes	For fibroscope and respiratory equipment
Orthophthaldehyde	Broad spectrum microbicidal and sporicidal	No activation required No fumes	Costly Stains equipment	For scopes Alternative to glutaraldehyde
Iodine compounds	Microbicidal spares M Tuberculosis and spores	Rapid action	Corrosive to metals, plastic and rubber Stains items	As an antiseptic
Alcohols	Wide microbicidal activity Non sporicidal	Non staining	Flammable	Hand disinfection For endoscopes
Phenols	Wide microbicidal activity Non sporicidal	Easily available Low cost	Irritant to skin Depigmentation	As surface disinfectant
Quaternary Ammonium compounds	Microbicidal spares M Tuberculosis, Not sporicidal, Not virucidal	Less irritant Good detergent property	Occupational asthma	As surface disinfectant For non-critical items
Peracetic acid	Broad spectrum microbicidal and sporicidal	No activation required Wide compatibility	Expensive Irritant to eye and skin	For fibroscopes
Chlorines	Wide microbicidal activity Non sporicidal	Low cost Fast acting	Corrosive to metals	Surface disinfectant To clean blood and body fluid spills
Hydrogen Peroxide	Broad spectrum microbicidal and sporicidal	No activation required	Serious eye damage Incompatible with some metals	Fogging of operating room For endoscopes
Formaldehyde	High level disinfectant	Non corrosive	Pungent odor and irritant fumes Carcinogenic	Withdrawn from use

3. Which of the following statements regarding reducing activation energy are true:

1. ΔGf is lowered
2. Stability of transition state is reduced
3. It depends upon complementarity of active site of enzyme and substrate
4. Conformational change may occur at the active site of enzyme at the time of binding to substrate

---

A. 1,2,3,4
B. 2,3,4
C. 3,4
D. 1,4


[TS]‡
ΔG‡uncat
E-[TS]‡
Uncatalyzed rxn
ΔG‡cat
Go
S
Catalyzed rxn
ΔG°rxn
P
Course of reaction
(reaction coordinate)

4. A 41-year-old woman is being evaluated for peripheral edema. The patient has gained 6.8 kg over the past 8 weeks. Her urine is "frothy." Blood pressure is 140/90 mm Hg and pulse is 80/min. Examination shows generalized edema. Heart sounds are normal. The abdomen is soft and nontender. Serum creatinine is 1.1 mg/dL. Urinalysis shows 4+ protein, 0-3 leukocytes/hpf, and oval fat bodies. Serum contains IgG4 antibodies to the phospholipase A2 receptor (PLA2R). Which of the following is the most likely diagnosis?

A. Focal segmental glomerulosclerosis
B. Membranous nephropathy
C. Minimal change disease
D. Mixed cryoglobulinemia

5. A 40-year-old nulligravida woman comes to the office with a 2-month history of worsening right pelvic pain. She experiences the pain daily, and its intensity does not vary during her menstrual cycle. In addition, the patient's waist size has increased despite a decreased appetite. Pelvic examination shows an irregularly shaped, fixed adnexal mass. One week later, she undergoes surgery to remove a large ovarian neoplasm. Within the true pelvis, the surgeon can most likely palpate the right ureter immediately anterior to which of the following structures?

A. Gonadal vein
B. Inferior vena cava
C. Internal iliac artery
D. Uterine artery

6. A 45-year-old male patient presents to their primary care physician with a complaint of dysphagia. He reports that he experiences a sensation of food getting stuck in his throat when he eats. He also reports some mild chest discomfort with exertion. The patient has a history of Down syndrome and congenital heart disease. Which imaging modality is typically used to diagnose dysphagia lusoria?

A. Chest X-ray
B. Electrocardiogram
C. Echocardiogram
D. Computed tomography angiography

7. A 35-year-old male, diagnosed with squint presented to the ophthalmology department for regular check-up. On examination, marked muscle was found affected. At which level does the nerve supply for the marked structure arise?

A. Red nucleus

B. Sub thalamic nuclei

C. Decussation of pyramidal tract

D. Olivary nucleus

8. A 10-year-old boy came with complaints of anaemia, weakness, with low weight to height ratio for his age profile. On examination, urine sample shows elevated level of orotic acid. Further study reveals low uridine monophosphate (UMP) synthase activity. What can be the best treatment to administer?

A. Adenine
B. Uridine
C. Hypoxanthine
D. Thymidine


PRPP
Glycine
De Novo
biosynthesis
AMP deaminase
GMP reductase
AMP
ADSL
AMPS
ADSS
IMP
IMPDH
XMP
GMPS
GMP
Inosine
Salvage synthesis
Hypoxanthine

9. A patient presented with fever, dyspnea, palpitations, and tachycardia. The echocardiography revealed vegetation in the septal leaflets. Tissue sections from them reveal the following findings as shown in the image. What is the most likely implicated agent?

A. Pigmented septate hyphae at 90 degrees, Mucormycosis

B. Right angle branching septate hyphae, Fusarium

C. Acute angle branching hyphae, Rhizopus

D. Acute angle branching septate hyphae, Aspergillus



10. 15-year-old boy comes to the OPD due to right ear itching and discomfort for the past several days. He has no fever or hearing loss but has noted scant drainage of thin, whitish fluid. The patient has been taking swimming lessons at a local gym. On examination, there is no redness around the ear, but gentle traction of the pinna elicits pain. During inspection of the external auditory canal, a speculum is inserted into the meatus in close contact with its posterior wall, causing the patient to suddenly become lightheaded and faint. He recovers spontaneously within a few minutes with no residual confusion. Which of the following nerves was most likely irritated during the procedure?

A. Facial nerve

B. Jacobson nerve

C. Arnold nerve

D. Auriculotemporal nerve

11. Identify the Diagram shown in the image below?



A. PERT management technique
B. Pie chart model
C. Epidemiological triad model
D. Sufficient component cause model

12. A 45-year-old woman is brought to the hospital after she collapsed during an airshow on a hot summer day. She is receiving amitriptyline for fibromyalgia. Temperature is 40.5 C (104.9 F), blood pressure is 90/60 mm Hg, pulse is 110/min, and respirations are 22/min. The skin is warm and red. She is disoriented. Neurologic examination shows no focal findings. If this patient's medication contributed to her current condition, which of the following mechanisms is most likely responsible?

A. Impaired cardiovascular response to heat stress

B. Impaired dissipation of body heat

C. Increased heat production by increasing muscle tone

D. Increased peripheral vasoconstriction

13. Identify the correct statement

1. Type I collagen is seen in bone and elastic cartilage

2. Reticular fibres are present in lymph nodes, bone marrow and thymus

3. Right-left movement is by atlanto-axial joint

4. Costo-transverse joints are plane synovial joints

A. 1,2,3

B. 2,3,4

C. 3,4

D. 1,2,3

14. Identify the false statement from the following:

A. True positive fistula test indicates perilymph fistula
B. False positive fistula test indicates congenital syphilis
C. True negative fistula test indicates late stages of Meniere’s disease
D. False negative fistula test indicates dead labyrinth

15. A 4-year-old girl is brought to the clinic due to an inward deviated right eye. Examination shows a pale red reflex in the right eye relative to the left eye. Dilated funduscopic examination reveals a well-circumscribed white mass within the retina. Genetic analysis shows a germline mutation in the patient's cells leading to development of the mass. Which of the following is the most likely function of the protein affected by this patient's gene mutation?

A. Activation of receptor tyrosine kinase

B. Conversion of GTP to GDP

C. Opposition of p53 activity

D. Prevention of the G1/S cell cycle transition

16. Identify the true statements:

1. **At physiological pH, arginine has a positive charge**
2. **Selenocysteine is coded by UGA**
3. **Heme synthesis occurs in BM, RBCS and hepatocytes**
4. **Vitamin C cannot be produced in humans due to lack of L-gulonolactone oxidase**

A. 1,2,3,4
B. 3,4
C. 1,2,4
D. 2,3

17. All of the following muscles have parallel oriented fibers except:

A. Sartorius
B. Rectus abdominis
C. Sternohyoid
D. Tibialis anterior

Muscles with parallel fasciculi

- Muscles in which fasciculi are parallel to the line of pull and have greater degree of movement.
- They are further classified based upon the shape:
  - *Quadrilateral*: Thyrohyoid
  - *Strap-like*: Sternohyoid and sartorius
  - *Strap-like with tendinous intersections*: Rectus abdominis
  - *Fusiform*: Biceps brachii, digastric

Muscles with oblique fasciculi

- Muscles fasciculi are arranged oblique to the line of pull. This increases the power of the muscle, but reduces the range of movement.
- They are further classified into:
  - *Triangular*: Temporalis, adductor longus, trapezius
  - *Unipennate*: Flexor pollicis longus, extensor digitorum longus
  - *Bipennate*: Rectus femoris, flexor hallucis longus
  - *Multipennate*: Tibialis anterior, subscapularis, deltoid (acromian fibers).

18. A 25-year-old homeless man with a history of schizophrenia is brought to the hospital by local police. He destroyed a television set at a local electronics store and then became extremely agitated and violent when employees attempted to intervene The patient's speech is difficult to follow, and he is distracted, seemingly listening to voices that only he can hear. He is admitted to the psychiatric ward and given haloperidol, which calms him down. Later that evening, however, he walks to the nurses' station and says, "What's happening to me?" The patient is upset and refuses to sit down, pointing to his neck. Physical examination shows a sustained contraction of his neck to the right side. Which of the following is the most appropriate next step in management?

A. Haloperidol

B. Diphenhydramine

C. Levodopa

D. Lorazepam

19. A 35-year-old, previously healthy man is evaluated immediately after accidental exposure to ionizing radiation at a fluoroscopy manufacturing facility. The patient is currently asymptomatic. Vital signs are normal. Physical examination shows no abnormalities. If radiation exposure is significant, which of the following cells are most likely to be affected first?

A. Bladder epithelial cells

B. Cardiac myocytes

C. Intestinal crypt cells

D. Renal proximal tubular cells

20. Match the following:

1. Stimulation of this area results in turning the eyes and head to the contralateral side
2. A lesion here results in nonfluent, telegraphic speech
3. Ablation in this area results in a contralateral upper homonymous quadrantanopia
4. A lesion here results in fluent speech with neologisms
5. A lesion of this area is characterized by finger agnosia, dyscalculia, dysgraphia, and dyslexia
6. Destruction of this area results in an aphasia characterized by fluent speech, good comprehension, and poor repetition
7. Lesions in this gyrus result in contralateral astereognosis
8. Lesions in this gyrus result in contralateral spasticity

A. 1-H, 2-F, 3-G, 4-D, 5-E, 6-C ,7-A, 8-B

B. 1-H, 2-G, 3-F, 4-E, 5-D, 6-C ,7-B, 8-A

C. 1-H, 2-G, 3-E, 4-F, 5-D, 6-B ,7-C, 8-A

D. 1-D, 2-G, 3-F, 4-E, 5-H, 6-C ,7-B, 8-A



21. A 5-month-old girl was brought to the casualty with seizures. The paediatrician tries to do CSF sampling. What are the structures punctured by the paediatrician while piercing through anterior fontanelle?

A. Scalp, dura, arachnoid

B. Scalp, epicranium, endocranium and dura

C. Scalp, synchondral membrane, dura, arachnoid

D. Pericranium, dura, arachnoid

22. A research group conducted a study to compare the levels of creatine kinase-MB (CK-MB) between type 2 diabetes mellitus (T2DM) patients given statin therapy. Participants were divided into 4 groups based on treatment. Groups I, I, and III consisted of T2DM patients who had been given statin therapy (atorvastatin, simvastatin, rosuvastatin, respectively) for at least 6 months. Group IV consisted of T2DM patients who had not been given statin therapy. Which of the following statistical tests is most appropriate to compare the CK-MB levels between Groups I, I, III, and IV in this study?

A. Analysis of variance

B. Chi-square test

C. Unpaired t-test

D. Paired t-test

23. Identify the condition showing the following variation in plasma proteins:

A. Multiple myeloma
B. Hepatic cirrhosis
C. Nephrotic syndrome
D. TB



24. Identify the incorrect pair:

1. **Talocalcaneonavicular joint – Saddle joint**
2. **Malleo-incudal joint-Ball and socket joint**
3. **Atlanto-axial joint-Pivot joint**
4. **Acromio-clavicular joint-Plane joint**

A. 1,3,4

B. 2,3,4

C. 1,2

D. 2,4

25. Identify the organism from the life cycle

A. Ancylostoma duodenale

B. Strongyloides stercoralis

C. Ascaris lumbricoides

D. Enterobius vermicularis



26. A 50-year-old man is scheduled to undergo a colonoscopy for routine colorectal cancer screening. He has no abdominal pain, gross blood in the stool, constipation, change in stool caliber, or unintentional weight loss. He also has no family history of colon cancer. The procedure is uncomplicated, but the patient is found to have a small mucosal lesion (~1 cm) in his sigmoid colon. The lesion is biopsied and its histology is shown in the image below. The pathogenesis of this patient's lesion most likely began with a mutation involving which of the following genes?

A. APC

B. STK11

C. SMAD4

D. MSH2





27. A 24-year-old male is brought to the psychiatry emergency department by his family members. They report that he has been muttering to himself, exhibiting suspicious behaviour, and believing that neighbours are conspiring against him by video recording him. He has not attended work for the last three months. Over the past month, he has become more withdrawn, speaking minimally, and has been observed standing in his room at the same place for hours. In the last two days, the patient has stopped speaking completely and has refused food and water. In the emergency department, the patient did not respond to any questions and stood as given in the image below for 30 minutes. What is the best next step in the management of this patient?

A. Risperidone
B. IV lorazepam
C. Electroconvulsive therapy (ECT)
D. Haloperidol

28. Which of the following is NOT one of the three clinical diagnostic criteria for bacterial vaginosis described by Amsel?

A. Abnormally high vaginal pH

B. Presence of an abnormal discharge and erythema of the vagina

C. Clue cells seen on vaginal saline preparation by light microscopy

D. Characteristic fishy odour release with addition of potassium hydroxide to vaginal secretions

29. Identify the incorrect pair

A. Montenegro test – Kala azar
B. Frenkel test – Malaria
C. Fairley test – Schistosomiasis
D. Bachman test – Trichinellosis

30. Match the following

1. Main elevator of mandible

2. Superior head originates from infratemporal surface and infratemporal crest of the greater wing

3. Receives supply from main trunk of trigeminal nerve

4. Originates from zygomatic arch and inserts on the angle of mandible

A. A-1, B-2, C-3, D-4

B. A-1, B-4, C-2, D-3

C. A-2, B-1, C-3, D-4

D. A-1, B-4, C-3, D-2



31. 35-year-old man with a history of bipolar and substance use disorders comes to the emergency department due to depression, auditory hallucinations, and suicidal ideation. His medications include lithium and escitalopram. The patient has a history of 5 psychiatric hospitalizations and 2 past suicide attempts, including overdose on his medications and attempted hanging. The patient has been using "anything I can get my hands on" because his depression is unbearable. He is hospitalized and placed on suicide precautions. His dose of escitalopram is increased to target his depression, and risperidone is added to treat the hallucinations. His lithium level is 1.0 mEq/L. On the second day of hospitalization, the patient reports muscle pains, abdominal cramping, nausea, and diarrhea. His temperature is 37.2 C (99 F), blood pressure is 130/85 mm Hg, and pulse is 84/min. The patient is alert and restless, and his pupils are dilated bilaterally. Bowel sounds are hyperactive and neurologic examination is normal. Which of the following is the most likely explanation for his symptoms?

A. Serotonin syndrome

B. Cocaine withdrawal

C. Lithium toxicity

D. Opioid withdrawal

32. Identify the true statements:

1. IVC passes through the central tendon of diaphragm
2. Left superior intercostal vein drains into the azygos vein
3. There are two dorsal interossei at the middle finger
4. Superior relation of epiploic foramina is formed by the hepatoduodenal ligament and right adrenal

---

A. 1,2,3,4
B. 1,2,3
C. 1,3
D. 2,4

33. Identify the false statements:
1) Rickettsial morula in a monocyte is diagnostic of Ehrlichia chaffeensis
2) E6, E7 are targets of vaccine against HPV
3) SARS CO-2 specimen handling for molecular testing needs BSL-2
4) Peptostreptococcus causes scromboid fish poisoning

A. 2, 4
B. 1,2,4
C. 3,4
D. 2,3

34. A 45-year-old previously healthy man comes to the office for evaluation of a neck lump. The patient first noticed the swelling a month ago while shaving, and it has progressively enlarged. On physical examination, there is a nontender, firm nodule in the upper portion of the left thyroid lobe. Laboratory studies reveal a normal serum TSH and elevated serum calcitonin levels. Neck ultrasonography shows a 2-cm hypoechoic thyroid nodule with microcalcifications. Fine-needle aspiration of the nodule reveals spindle-shaped cells in an amorphous background. Which of the following gene alterations most likely contributed to this patient's current condition?

A. Activation of ALK

B. Activation of RET

C. Inactivation of MEN1

D. Activation of GNAS

35. Identify the distribution represented by ‘diagram 3’ in the Image below?

A. Normal distribution
B. Normal distribution with outliers
C. Positive skew
D. Negative skew



36. If changes made in some of the sequence of amino acids in the chain of human proinsulin, which of the following changes will not make any difference in the biochemical activity of the molecule?

A. Breaking disulfide linkage
B. Interchange of A1 and A4
C. Interchange of B29 and B30
D. Interchange of A5 and A6



37. Which procedure is shown in the following picture?



A. Infratrochlear block
B. Mandibular nerve block
C. Anterior ethmoidal block
D. Maxillary nerve block


Infratrochlear nerve
Supratrochlear nerve
External branch of anterior ethmoidal nerve
Infraorbital nerve

38. Match the following with the appropriate diagnosis:

A. Lichen planus
B. Anti-synthetase syndrome
C. Osteoarthritis
D. Dermatomyositis

A. 1-a, 2-d, 3-c, 4-b
B. 1-a, 2-d, 3-b, 4-c
C. 1-d, 2-a, 3-c, 4-b
D. 1-d, 2-a, 3-c, 4-d

39. The nerve being tested here is injured most likely at what site?

A. Between two heads of pronator teres
B. At the level of supinator muscle
C. Radial groove
D. Medial epicondyle



40. Mr. Ramesh, a 55-year-old male patient, has been diagnosed with laryngeal malignancy and is scheduled to undergo modified radical neck dissection (MRND). The surgery is being performed in a sitting position to provide the surgeon with better access to the operative field. The anaesthetic team has taken all necessary precautions to prevent any complications during the surgery. During the surgery, while ligating the internal jugular vein, the patient's end-tidal carbon dioxide (etCO2) dropped from 40 mmHg to 14 mmHg. The patient suddenly developed hypotension along with cardiac arrhythmia, and the surgical team immediately recognized the emergency situation. Which of the following is the most likely cause?

A. Sympathetic over activity

B. Vagal stimulation

C. Venous air embolism

D. Carotid body stimulation

41. In which of the given cases is this procedure usually performed?

A. MRM in node negative patients

B. MRM in node positive patients

C. BCT in node negative patients

D. BCT in node positive patients



42. A 66-year-old woman with warfarin induced bleeding receives IV vitamin K, 4 U PRBCs and 4U FFP. 4h later, she begins to complain of shortness of breath. Vitals T - 39°C, RR 35/min, SpO2 84% on RA, and BP 88/52 mm Hg. Exam revealed B/L crackles and the patient was intubated for respiratory failure. CXR is as shown in the picture. What is the likely diagnosis?

A. Anaphylaxis

B. TACO

C. TRALI

D. Aspiration



43. Which of the following is not a correct match with respect to the mechanism of action and clinical use of the given drugs?

A. Zavegepant - CGRP receptor antagonist used for the treatment of acute migraine attacks

B. Sparsentan - Dual endothelin and angiotensin receptor antagonist used for the treatment of CHF

C. Lenacapavir - HIV-1 capsid inhibitor used for the treatment of HIV infection

D. Teplizumab - Anti-CD3 monoclonal antibody used for the prevention of type 1 diabetes in high-risk individuals

NEW DRUGS

- Daprodustat: HIF-PH inhibitor: Anemia due to chronic kidney disease
- Elacestrant: SERD: Hormone receptor-positive breast cancer
- Tirzepatide: Dual GLP-1 and GIP receptor agonist-Type 2 diabetes
- Mavacamten: Cardiac myosin inhibitor-HOCM

44. You are preparing to intubate a patient in the operating room. As part of your preoperative assessment, you perform a Mallampati examination to assess the patient's airway. Identify the class from the image shown below?

A. Mallampati Class I

B. Mallampati Class II

C. Mallampati Class III

D. Mallampati Class IV

45. A 4-year-old boy is brought to the emergency department for difficulty breathing and a productive cough. He has a past medical history of recurrent episodes of sinusitis and otitis media. The respiratory infections typically have a protracted course but respond to high-dose antibiotic therapy. Pulmonary auscultation shows crackles and wheezing. Cardiovascular examination shows a cardiac point of maximal impulse that is palpated at the right 5th intercostal space. Which of the following is most likely to be dysfunctional in this patient?

A. C1 complement component

B. Chloride channels

C. Dynein arms

D. IL-2 receptors

46. Correct about Brown-Sequard syndrome:

A. Extensive loss of pain on same side below the level of hemisection

B. Loss of fine touch and vibration on opposite side below the level of lesion

C. Extensive paralysis of muscle on opposite side below the level of lesion

D. Complete LMN type paralysis on same side at the level of lesion

47. A 2-year-child presented with complaints of abdominal pain following which he develops altered sensorium. He lives in an old house with paint chipping off. The following appearance is noted on X-rays. Which enzyme deficiency is likely in this child causing all these symptoms?

A. ALA synthase
B. Heme synthase
C. ALA dehydratase
D. Coproporphyrinogen synthase



48. All are true about the muscle whose function is tested in the below shown test except:

A. Inserts into the lesser tubercle of the humerus
B. Primary function is external rotation
C. Arises from the subscapularis fossa
D. Supplied by a branch of the posterior cord of the brachial plexus

49. A 45-year-old male with a history of human immunodeficiency virus (HIV) presents to the clinic with complaints of fatigue, weakness, and shortness of breath. The patient reports taking his anti-HIV medications consistently, which includes zidovudine (AZT). A complete blood count reveals a hemoglobin level of 8.5 g/dL, and a peripheral blood smear shows macrocytic anemia. Which of the following adverse effects of zidovudine is most likely causing the patient's symptoms?

A. Lactic acidosis
B. Peripheral neuropathy
C. Pancreatitis
D. Bone marrow suppression

50. You are the team leader during a cardiac arrest resuscitation scenario in a hospital setting. A 65-year-old patient has suddenly collapsed and is unresponsive. The monitor displays a rhythm as shown in the image. What is the next immediate step?

A. Administer epinephrine

B. Begin high-quality CPR

C. Defibrillate at 360 joules

D. Administer $MgSO_4$



51. A 24 year old G2 at 28 weeks is brought to the emergency after being found unconscious at home by her mother. She is disoriented on arrival and her BP is 180/110 mmHg and 4 + proteinuria. On arrival at the emergency, she has a tonic-clonic seizure which resolves with MgSO4 administration. Once the patient is stabilized, she undergoes cesarean delivery for fetal distress. Post-partum, the below MRI is done. What is the most likely finding on imaging?

A. Subdural hematoma
B. Parenchymal hemorrhage
C. Subarachnoid hemorrhage
D. Posterior reversible encephalopathy syndrome



52. A screening test is positive in 50% diseased population and 10% healthy population. Calculate the PPV:

A. 83

B. 90

C. 50

D. 89

53. A 63-year woman old presents to the AIIMS emergency chest pain after an RTA. Which resulted in crushed left lower limb. ECG shows 1-2 mm ST elevations in leads V2-V4 & troponin are elevated. The patient undergoes CAG which was normal. Which of the following is likely?

A. Aortic Dissection

B. Ramus intermedius

C. Severe mitral regurgitation with papillary muscle dysfunction

D. Akinetic LV apex with ballooning

54. 6-year-old girl is brought to the office for evaluation of short stature. Her parents have noticed that she bears little resemblance to her 2 older siblings. The patient's mother and father are 172.5 cm (5'8") and 183 cm (6') tall, respectively. On physical examination, the patient's height is 3 standard deviations below the mean for her age. Other findings include low-set ears, a high arched palate a webbed neck, and cubitus valgus. Chromosomal analysis reveals a 45,XO karyotype. The patient is started on a medication to improve growth and normalize her height. Which of the following intracellular pathways is stimulated by the medication used in this patient?

A. Binding of activated receptors to DNA to modify transcription

B. Cyclic AMP-Protein kinase A pathway

C. Inositol triphosphate calcium-calmodulin pathway

D. JAK-STAT pathway

55. A 40-year-old woman has noted painless enlargement of her neck for the past month. On physical examination, she has diffuse, symmetrical thyroid enlargement without tenderness on palpation. Laboratory studies show her serum TSH is elevated but serum thyroxine is lower side of normal limits. Her microscopic examination is shown here. Which of the following laboratory test findings is she most likely to have?

A. Increased serum calcitonin
B. High anti-thyroid peroxidase antibody titre
C. Increased serum parathormone
D. Positive antinuclear antibody test



56. A 23-year-old medical student presents to the surgery emergency department after cutting her wrist with a blade. After receiving first aid, the resident notices multiple healed cut marks on her wrist. Upon psychiatry referral, the patient informs the resident that she cut herself because she was feeling very angry at her boyfriend who spent the entire evening watching an IPL match rather than calling her. She also mentions that she has never been in a relationship with a guy for more than 2 months and believes that all men are horrible, while women are the best. What is the most likely diagnosis for this patient?

A. Histrionic Personality Disorder
B. Borderline Personality Disorder
C. Narcissistic Personality Disorder
D. Dependent Personality Disorder

57. All are true about the part labelled here except:

A. Derived from a type of atavistic epiphyses
B. Gives insertion to pectoralis major muscle
C. Short head of biceps takes origin from here
D. Can be felt in the infraclavicular fossa on examination



58. Identify the correct statements:

1. Grade 2 disability per 10 lac population is the most important leprosy awareness indicator

2. VVM shows cumulative exposure of the vaccine to the heat and assess vaccine potency

3. As per the National Tuberculosis Elimination Program (NTEP), HIV-TB prevalence of district >15% is a high-priority district

4. The external validity of a screening test is related to generalizability

Options:

A. 1,2,3,4

B. 1,2,4

C. 1,2,3

D. 2,3,4

59. A7-year-old girl is brought to the office due to a diffuse rash. The father says she developed a lace-like, erythematous rash this morning on her stomach, back, and extremities. Two days ago, the girl developed red, flushed cheeks. She also had congestion, headache, and a low-grade fever for 3 days earlier this week. The infectious agent most likely responsible for this patient's disease replicates in progenitors of which of the following cell lines?

A. Erythrocytes

B. Hepatocytes

C. Intestinal epithelial cells

D. Lymphocytes

60. Anaerobic glycolysis of which of these produces 3 ATPs perunit glucose consumed?

A. Glucose
B. Fructose
C. Galactose
D. Glycogen

61. 44-year-old man comes to the OPD for follow-up of type 2 diabetes mellitus. He was diagnosed with diabetes 4 years ago and has been taking metformin since then. Six months ago, a new antidiabetic agent was added due to suboptimal glycemic control. Today, the patient is happy because he has lost 7 kg without any changes in diet or activity since the new therapy was started. His only symptoms are occasional nausea vomiting, and a prolonged sensation of fullness after eating. Laboratory testing reveals normal renal function studies and hemoglobin A1c of 7.2%. If this patient's weight loss is due to the new medication, which of the following agents is most likely responsible?

A. Basal-bolus insulin therapy
B. Thiazolidinediones
C. Glucagon-like peptide-1 agonist
D. Sulfonylurea

62. Identify the correct pair

1. **Benefit measured in terms of natural units-Cost benefit analysis**
2. **Method based on behavioral sciences-MBO**
3. **PERT-Network Analysis**
4. **Benefit measured in terms of QALY-Cost effective analysis**

Options;

A. 1,2,3

B. 2,3,4

C. 2,3

D. 1,4

63. A 20-year-old woman is brought to the emergency department by police at 2:30 AM after she was caught attempting to scale a fence at the Rashtrapati Bhawan. The patient appears highly agitated and paces around the examination room. She has just flown in from out of state to "meet with the president about a foolproof plan for eliminating worldwide terrorism." The patient has barely slept for the past week due to her intensive work on this plan. The evaluation has to be stopped when the patient begins banging on the door and demanding to leave. Temperature is 37 C (98.6 F), blood pressure is 148/84 mm Hg, pulse is 98/min, and respirations are 22/min. Urine drug screen is negative. Administration of which of the following medications is the most appropriate next step in management of this patient?

A. Lithium

B. Clozapine

C. Haloperidol

D. Olanzapine

64. A newborn is being evaluated in the nursery. The patient was delivered to a 23-year-old woman. The mother developed dark terminal facial hair and a deepened voice during the pregnancy. The delivery was unremarkable, and the patient's vital signs are within normal limits. Examination shows ambiguous genitalia land clitoromegaly. Laboratory studies reveal elevated serum levels of testosterone and androstenedione. Karyotype testing shows a 46,XX genotype. Ultrasound of the pelvis shows a normal-sized uterus. This infant's presentation is most likely due to deficiency of which of the following enzymes?

A. 5-Alpha reductase

B. 17-Alpha hydroxylase

C. 21-Hydroxylase

D. Aromatase

65. Which of the following is included in high-quality CPR?

1. **Compression depth of at least 5 cm**
2. **Rotate chest compressor every 3 mins**
3. **Avoid excessive ventilation**
4. **Compression - ventilation ratio of 30:2**
5. **Allow complete recoil**
6. **ET co2 <10mm**
7. **Intra-arterial DBP >20mm Hg**
8. **Breaths 20-30/min**

Options;

A. 1,2,3,4,5,6,7,8

B. 1,2,4,5,7,8

C. 1,3,4,5,7,8

D. 1,3,4,5,7

66. 38-year-old woman, gravida para 0, at 34 weeks gestation comes to the OPD due to increasing swelling in her hands and legs. The patient first noticed the swelling a few weeks ago but now cannot stand for more than an hour without needing to elevate her legs. She has no orthopnea, dyspnea, or chest pain. The patient has no chronic medical conditions and takes no daily medications. Vital signs are normal. Cardiac examination is within normal limits. Lungs are clear to auscultation. Bilateral lower extremities have 2+ pitting edema to the midshin. Urine dipstick shows no protein. Compared to a healthy nonpregnant woman, which of the following parameters is most likely to be decreased in this patient?

A. Cardiac preload

B. Cardiac output

C. Central venous pressure

D. Systemic vascular resistance

67. A 19-year-old man comes to the physician due to diarrhea for the last 4 months. He visited a student health clinic at his university a month ago and was prescribed a course of antibiotics, but his diarrhea failed to improve. He also has persistent nausea, a bloating sensation, and a 6.8-kg weight loss. Microscopic examination of his stool is shown in the image. Impairment of which of the following immune functions would most likely predispose to this patient's infection?

A. Bacterial killing by neutrophils

B. CD8+ T lymphocyte-mediated cytotoxicity

C. Eosinophil-mediated cytotoxicity

D. Secretory IgA production

68. All of the following structures lie in the relation to the left ureter except:

A. Mesentery of sigmoid colon
B. Bifurcation of common iliac artery
C. Quadratus lumborum
D. Gonadal vessels

Relations of Ureter

Anterior Relations

Right ureter

- Duodenum (2nd part)
- Right colic vessels
- Ileocolic vessels
- Right gonadal vessels
- Root of mesentery.

Left ureter

- Left colic vessels
- Sigmoidal vessels
- Left testicular or ovarian vessels
- Sigmoid mesocolon.

Posterior Relations

Both ureters run anterior to psoas major muscle and bifurcation of common iliac artery.

69. 65-year-old man presents with progressive dyspnoea & pulmonary function testing was done. FEV1 is 2.5 L (78% predicted), FVC is 4.00 L (94% predicted), and the FEV1/FVC ratio is 62.5%. The flow-volume curve is shown. What is the most likely cause?

A. Aspirated foreign body
B. U/L vocal cord paralysis
C. Subglottic stenosis
D. COPD



70. A 4-month-old girl is diagnosed with a febrile urinary tract infection. A voiding cystourethrogram is shown in the image. Which of the following is the most likely cause of this infant's presentation?

A. Congenital short intravesical ureter length

B. Damage to the rectovaginal septum

C. Obstruction due to a renal stone

D. Presence of a posterior urethral valve

71. Identify the correct kit for the clinical presentations:

1. Painful ulcers upon palpation, with suppurative inguinal lymphadenopathy on the right side

2. Vaginal discharge in a CSW with cervical erosions

3. Painful, swollen, and tender inguinal lymph node in a truck driver

4. Painless ulcer in the genital area with non-tender inguinal lymphadenopathy in CSW; past h/o treatment of a similar lesion resulted in fever, malaise, tachycardia and vomiting

---

Options;

A. 1-Red, 2-Green, 3-Black, 4-Blue

B. 1-White, 2-Green, 3-Black, 4-Blue

C. 1-White, 2-Grey, 3-Black, 4-White

D. 1-Blue, 2-Grey, 3-White, 4-Blue

72. A 24y P2L2 who does not want more children presents to the OPD as she cannot feel the thread of the Multiload 375 that was inserted 3 years earlier. Her last menstrual period was 3 weeks earlier. On examination, the cervix appears normal, and the thread is not seen. A transvaginal scan shows the following image. What is the next best step?

A. Hysteroscopic IUCD removal
B. Reassure her
C. Remove the IUCD with a hook
D. X-Ray Abdomen and Pelvis



73. A 45-year-old male presents with fatigue, night sweats, and weight loss. A complete blood count (CBC) is performed, and the peripheral blood smear shows the presence of cells with increased basophilic cytoplasm, granulocytic precursors, and a high percentage of blasts. A bone marrow biopsy is performed. Which of the following is the most appropriate investigation for this patient's condition?

A. Flow cytometry
B. Cytogenetics
C. FISH
D. Karyotyping



74. 37-year-old woman, gravida 2 para 1, at 30 weeks gestation comes to the OPD for a routine prenatal examination. Her pregnancy has been uncomplicated to date, and fetal movement is present. At the last visit, she drank a solution containing 50 g of glucose, and serum glucose was abnormally elevated an hour later. Which of the following hormones is most likely contributing to this patient's abnormal laboratory findings?

A. HCG
B. Human placental lactogen
C. Oxytocin
D. GH

75. 31-year-old woman comes to the office due to transient visual changes that are characterized by "loss of vision for a minute" and commonly precipitated by bending forward or lifting objects. During the last 3 months, she has had persistent headaches that she attributes to migraines. The patient takes over-the-counter analgesics as needed. Blood pressure is 140/90 mm Hg and pulse is 72/min and regular. BMI is 32.4 kg/m2. Funduscopic findings of the left eye are shown in the image below, with similar findings in the right eye. Which of the following is the most likely cause of the observed funduscopic findings in this patient?

A. Choroidal inflammation
B. Elevated intracranial pressure
C. Elevated intraocular pressure
D. Optic nerve inflammation

76. A 25-year-old woman comes to the emergency department due to 2 days of abdominal pain and bloody diarrhea. Temperature is 37.1 C (98.8 F), blood pressure is 119/76 mm Hg, pulse is 92/min, and respirations are 16/min. There is abdominal tenderness with no rebound or guarding. Guaiac-positive bloody stools are detected on rectal examination. Stool cultures reveal an Escherichia coli strain that does not produce glucuronidase and does not ferment sorbitol on sorbitol-containing MacConkey agar. Which of the following best describes the mechanism of action of the toxin specific to these bacteria?

A. Disruption of the cellular cytoskeleton

B. Inactivation of 60S ribosomal subunits

C. Inactivation of elongation factor 2

D. Inhibition of sodium reabsorption

77. Identify the true statements:

1. **TxA2 has platelet aggregating and vasoconstricting effect**
2. **Lipoxin B4 inhibits inhibit neutrophil chemotaxis and adhesion to endothelium.**
3. **Il-1, Il-18, TNF are pyrogenic cytokines**
4. **C3 can be activated is both common as well as alternative pathways**
5. **T cells with a TCR composed of a γ and δ subunit are found associated with GI mucosa**

Options;

A. 1, 2, 3, 4

B. 1, 2, 4, 5

C. 2, 3, 4, 5

D. 2, 4

78. 65-year-old woman is concerned about her risk of fracture as her mother was hospitalized for osteoporotic hip fracture in the past. She has a history of hypertension, hyperlipidemia, coronary artery disease, seizure disorder, and gastroesophageal reflux disease. The patient underwent menopause at age 52. She has smoked a pack of cigarettes daily for 24 years but does not drink alcohol. Weight is 56 kg. Long-term use of which of the following medications may increase this patient's risk of osteoporosis and hip fracture?

A. Atorvastatin

B. Levetiracetam

C. Metoprolol

D. Omeprazole

79. Identify the nerve block given at the point shown:

A. Pudendal nerve
B. Sciatic nerve
C. Obturator nerve
D. Inferior rectal nerve



80. 7-year-old boy is brought to the clinic for a routine health maintenance examination. He underwent a splenectomy a year ago for persistent anemia and jaundice. He has received the pneumococcal and meningococcal vaccinations and takes penicillin prophylaxis. Physical examination reveals a well-appearing child. The sclerae are anicteric. The abdomen is soft and nontender; no masses are palpable. Several well-healed laparoscopy scars are noted. Complete blood count is as follows:
Hemoglobin: 11.3 g/dL
Reticulocytes: 3%
Platelets: 160,000/mm³
Leukocytes: 8,200/mm³
Peripheral blood smear with Wright stain is shown.
This patient's findings are most likely related to which of the following?

A. Chronic Hb oxidation and precipitation
B. Inability to produce beta globin chains
C. Mechanical red-blood-cell membrane damage
D. Persistence of red blood cell precursor nuclei



81. A 30 year old man presented with profuse bleeding from the nose which did not stop on applying local pressure or vasoconstrictors. Anterior nasal packing was done but the bleeding could still not be controlled. It was decided to embolize one of the arteries supplying the vascular plexus responsible for most of the anterior nasal bleeds. Which of the following arteries should never be embolized?

A. Sphenopalatine artery
B. Septal branch of superior labial artery
C. Anterior ethmoidal artery
D. Greater palatine artery

82. A 52-year-old woman comes to the emergency department due to recurrent right lower extremity swelling, pain, and erythema. The patient's medical history includes type 2 diabetes mellitus and depression. Her medications include metformin, paroxetine, and a multivitamin. The patient is diagnosed with cellulitis, admitted to the hospital, and started on antibiotics. She is continued on her home medications. Three days later, the patient becomes agitated and delirious with severe abdominal cramps and diarrhea. Her temperature is 39.2 C (102.6 F). blood pressure is 180/100 mm Hg, and heart rate is 120/min and regular. On examination, the patient is diaphoretic and tremulous, and her pupils are dilated. Bilateral hyperreflexia and ankle clonus are present. She begins to have seizures. Which of the following antibiotics was the patient most likely given to treat the cellulitis?

A. Clindamycin
B. Doxycycline
C. Linezolid
D. Vancomycin

83. Which of these is an example of anaplerotic reaction?

A. Pyruvate to Acetyl CoA
B. Pyruvate to oxaloacetate
C. Pyruvate to lactic acid
D. Pyruvate to acetaldehyde

84. Match the following drugs with their specific side effects:



A. Emtricitabine
B. Abacavir
C. Rosiglitazone
D. Liraglutide
E. Canagliflozin
F. Dabigatran
G. Warfarin

Options;

A. 1-B, 2-C, 3-F, 4-G

B. 1-A, 2-C, 3-E, 4-G

C. 1-G, 2-F, 3-E, 4-A

D. 1-E, 2-B, 3-C, 4-D

85. Identify the incorrect statements:

1. Superficial anterior compartment of leg has no neurovascular bundle

2. Tectorial membrane is the superior continuation of PLL

3. After parotidectomy, a person complains of loss of sensation in the shaving area. This is likely due to injury to auriculotemporal nerve.

4. Common carotid artery divides at the level of lower border of thyroid cartilage

5. Ansa cervicalis lies on the posterior wall of carotid sheath

---

A. 1,2,3,4,5

B. 1,4

C. 2,4,5

D. 1,3,4,5

86. A 70-year-old man, a retired banking executive with a history of smoking is admitted in ward with shortness of breath. Interpret the acid base disorder based on the arterial blood gas analysis report mentioned below. pH – 7.30, PCO2 -65 mm Hg, PO2 -60 mmHg, HCO3 - 36 mEq/L.

A. Metabolic acidosis
B. Respiratory acidosis
C. Metabolic alkalosis
D. Respiratory alkalosis

87. A medical student have given a topic to present in class. The topic given to you was excitation-contraction coupling. It is a process in which muscle cells are stimulated to contract and it is followed by muscle cell relaxation. It involves the following events:

1. **Action potential**
2. **Calcium release into the cytoplasm**
3. **Cross-bridge cycling**
4. **Cytoplasmic calcium removal**
5. **Depolarization of T tubules**

Which of the following options lists the correct sequence of events for excitation-contraction coupling in skeletal muscle tissue?

Options:

A. 1, 5, 2, 3, 4

B. 1, 2, 3, 4, 5

C. 1, 3, 4, 2, 5

D. 1, 4, 2, 5, 3

88. A female presents to the emergency 2 weeks after a normal vaginal delivery with abnormal involuntary flexure of the neck towards one side. History taking revealed that she was unable to lactate spontaneously for which she took a medication that she found on the internet without taking proper medical advice. Intake of which of the following drugs is most likely to lead to this condition?

A. Bromocriptine
B. Levodopa
C. Chlorpromazine
D. Benzhexol

89. All of the following are true:

1. **Leptin acts via JAK-STAT signalling pathway**
2. **Damage to the Zona Occludens can result in increased permeability in Vibrio cholera diarrhea**
3. **Raised 2,3 BPG, Acidosis, HbF cause a right shift of the oxygen-hemoglobin dissociation curve**
4. **Macula densa is located at the Junction of thick ascending limb and DCT**
5. **The organ with the least arteriovenous pressure difference is the kidney.**

Options;

A. 1, 2, 3, 4, 5

B. 2, 3, 4

C. 1, 2, 4, 5

D. 5

90. A 30-year-old female came to the OPD with complaints of backache. On examination, her eyes and pinna revealed blackish discoloration. What is the likely diagnosis?

A. DISH
B. Fluorosis
C. Alkaptonuria
D. Ankylosing spondylitis

91. A 15-year-old male presents to emergency department in a comatose state. He was found in this state by his friend in the schoolyard. His friend confesses that the patient has been depressed over the past several days. His initial labs reveal hypocalcaemia and a severe metabolic acidosis. The urine shows calcium oxalate crystals. What was the toxin ingested?

A. Methanol

B. Ethylene Glycol

C. Organophosphates

D. Ethanol

92. A 64-year-old man is evaluated for persistent fever and weakness. He has a history of aortic valve replacement for aortic stenosis. Physical examination reveals a new cardiac murmur with scattered petechia and splinter hemorrhages seen on his extremities. Echocardiogram shows a vegetation involving one of the aortic valve leaflets, and blood cultures grow enterococci. As part of the patient's treatment, 240 mg of intravenous gentamicin is started. The pharmacy calculates that, in this patient, gentamicin has a volume of distribution of 30L, a half-life of 4 hours, and demonstrates first-order kinetics. Which of the following is the most likely serum drug concentration just before the next dose 8 hours later?

A. 0.5 mg/L

B. 1 mg/L

C. 1.5 mg/L

D. 2 mg/L

93. Identify the correct statements

1. KEYNOTE-189 trial investigated NSCLC treatment using this drug that acts via pathway B.

2. Drug acting via pathway B has a role in treatment of ca endometrium.

3. Osimertinib is specifically used for treating NSCLC patients with T790M mutations and acts via pathway A.

4. Ipilimumab is a medication used in the management and treatment of metastatic melanoma that acta via pathway C.

A. 1, 2, 3, 4

B. 2, 4

C. 1, 2

D. 3, 4



94. What is the mechanism of action of the toxin produced by the bacterium shown in the figure below?

A. Increase in cAMP
B. Increase in cGMP
C. ADP-ribosylation of EF2
D. ADP-ribosylation of G protein



95. A neonate is lethargic, drowsy, and irritable with blood analysis showing hyperammonemia and orotic aciduria. Considering this patient, orotic acid levels are high due to which of the following?

A. Elevated ammonia
B. Elevated glutamine
C. Bypassing carbamoyl phosphate synthetase II (CPS-II)
D. Bypassing aspartate transcarbamoylase


Mitochondria
CO_2 + NH_3
Carbamoyl phosphate synthetase 1 (CPS-1)
2ATP
2ADP
Carbamoyl phosphate
Ornithine transcarbamoylase (OTC)
Ornithine
Citrulline
OTC Deficiency and its effects
Glutamine + 2ATP + CO_2
CPS-II
Carbamoyl phosphate
Carbomoyl Aspartate
Few steps
Orotic acid
Oritidine monophosphate (OMP)
Uridine Monophosphate (UMP)
Pyrimidines

96. Which of the following statements is/are true regarding anti-tubercular drugs?
1. Isoniazid inhibits mycolic acid synthesis in the cell wall of Mycobacterium tuberculosis.
2. Rifampicin is primarily used for the treatment of active tuberculosis and can cause liver toxicity and flu-like symptoms.
3. Bedaquiline is a second-line anti-tubercular drug that can cause QT prolongation and requires cardiac monitoring.
4. Pretomanid is primarily used as a monotherapy for the treatment of tuberculosis.

A. All options are correct
B. 1, 2 and 3 are correct
C. 1, 3 and 4 are correct
D. Only 1 and 2 are correct

97. A 42-year-old man is admitted to the hospital after sustaining fractures of his right femur and tibia in a motor vehicle collision. The patient becomes agitated, hypervigilant, and paranoid 12 hours after admission. He repeatedly tells the nurses that he can hear people in the corridor making insulting remarks about him. When hospital staff check the area near the patient's room, no one is found. Medical history is unremarkable, but psychiatric history is significant for cocaine, marijuana, and alcohol misuse. On examination, the patient is alert and oriented. Temperature is 37.2 C (99 F), blood pressure is 135/87 mm Hg pulse is 85/min, and respirations are 18/min. He is slightly tremulous and diaphoretic. Which of the following is the most likely diagnosis in this patient?

A. Alcoholic hallucinosis
B. Brief psychotic disorder
C. Cocaine withdrawal
D. Delirium tremens

98. The cranial nerve which is a content of the space labelled in the image below passes through which cranial foramina:

A. A
B. B
C. C
D. D



99. 64-year-old man comes to the emergency department due to 2 days of fever, chills, productive cough. The patient has a history of hypertension, hyperlipidemia, and type 2 diabetes mellitus. Oxygen saturation is 94% on room air when lying on his right side but drops to 89% when he lies on his left. Dullness to percussion and bronchial breath sounds are present on the left side. Imaging is shown here. Which of the following pathophysiologic mechanisms in the left lung is most responsible for the drop in oxygen saturation after the change in position?



A. Dead space ventilation
B. Effort dependent hypoventilation
C. Hypoxic vasoconstriction
D. Intrapulmonary shunting

100. 46-year-old man is admitted to the hospital for elective cholecystectomy. He is started on intravenous fluids and antibiotics. Just before he is to go to an operating room, the patient complains of a sudden onset of chest tightness and light-headedness. His medical history is significant for anxiety, and he has had 2 previous visits to the emergency department for panic attacks. His blood pressure is 118/67 mm Hg. His electrocardiogram is shown here. Which of the following is the best initial step in treating this patient?



A. Adenosine
B. Alprazolam
C. Amiodarone
D. DC Cardioversion

101. Glycogen synthesis and breakdown takes place in the same cell, having enzymes necessary for both the pathways. Why the glucose-6-phosphate, freshly synthesized during glycogenesis in cytoplasm of hepatocytes, is not immediately degraded by the enzyme glucose-6-phosphatase?

A. The thermodynamics does not favor such a reaction to occur
B. Glucose-6-phosphatase is present in the endoplasmic reticulum and cannot act on glycogen formed in the cytoplasm
C. Glycogenesis and glycogenolysis are tightly regulated such that enzymes of only one of those are present at a time.
D. Steric hindrance due to albumin

102. A patient has the following laboratory findings:
Sodium – 138 mEq/L
Potassium – 4.4 mEq/L
Chloride – 112 mEq/L
Bicarbonate – 8.0 mEq/L
Arterial pH – 7.21
Which of the following is the most likely acid-base status?

A. Anion gap metabolic acidosis
B. Anion gap metabolic acidosis plus metabolic alkalosis
C. Anion gap metabolic acidosis plus normal anion gap metabolic acidosis
D. Normal anion gap metabolic acidosis

103. Healthy adult volunteers are enrolled in a phase I clinical trial investigating the properties of a newly developed oral antimicrobial agent. The drug is administered in different amounts to the volunteers over the course of several weeks to determine the best dosage that minimizes toxicity while maintaining trough levels above the minimum inhibitory concentration. While reviewing the data, the researchers note that the drug's half-life seems to vary amongst the study participants. An increase in which of the following pharmacologic parameters is most likely responsible for the longer half-life seen in certain individuals?

A. Drug glucuronidation

B. Glomerular filtration rate

C. Oral bioavailability

D. Volume of distribution

104. Identify the correct statements with respect to the deficits in different brain regions:

1. **Lateral geniculate body-Keyhole shaped defect**
2. **Substantia nigra-Paralysis agitans**
3. **Putamen-Athetosis**
4. **Hypothalamus-Hemiballismus**

A. 1,2,3

B. 2,3,4

C. 1,2

D. 3,4

105. A person presents with pneumonia. His sputum was sent for culture. The bacterium obtained showed gram positive cocci in chains on microscopy and alpha haemolytic colonies on sheep agar. Which of the following will help in confirming the diagnosis?

A. Novobiocin
B. Optochin
C. Bacitracin
D. Oxacillin

106. A 30y P1 with a last menstrual period 10 days ago presents to the emergency with complaints of worsening right pelvic pain. She says that she had sudden onset of sharp pain after bending over to pick up her child. The pain did not respond to acetaminophen and has intermittently worsened during the past several hours. She also had 2 episodes of vomiting. On examination, there is abdominal guarding. Transvaginal sonography demonstrates an 8-cm enlarged right ovary. There is high impedance arterial flow of the ovary on colour Doppler. These findings are most consistent with which of the following clinical diagnoses?

A. Appendicitis
B. Ovarian torsion
C. Haemorrhagic cyst
D. Ruptured corpus luteum cyst

107. Identify the true statements

1. In amyloidosis, Beta pleated sheet will be seen in X-ray crystallography.

2. There is decrease in thyroid-binding globulin, transferrin, and fibrinogen in nephrotic syndrome.

3. Blood transfusion requires screening of donor blood for HIV, HCV, HBV but not dengue.

4. Massive splenomegaly is noted in hairy cell leukaemia, CML and aplastic anemia.

A. 1, 2, 3, 4

B. 1, 3

C. 1, 2, 3

D. 2, 4

108. 46-year-old man comes to the OPD due to right shoulder pain. The pain started 2 months ago after the patient began working at a new warehouse where he frequently moves heavy boxes. On physical examination, external rotation of the right shoulder against resistance is painful, but there is no weakness. MRI reveals near complete tear of the right infraspinatus muscle. The absence of weakness in this patient is best explained by compensatory hypertrophy of which of the following muscles?

A. Latissimus-dorsi
B. Subscapularis
C. Supraspinatus
D. Teres minor

110. Match the following with the mode of cellular transport:

1. Oxygen	A. Simple diffusion
2. GLUT	B. Facilitated diffusion
3. SGLT	C. Primary active transport
4. Na+/Iodide symporter	D. Secondary active transport
5. Na+/K+ ATPase	
6. Proton pump	

Options:

A. 1-A, 2-B, 3-B, 4-C, 5- D, 6-D

B. 1-B, 2-A, 3-B, 4-C, 5- D, 6-D

C. 1-A, 2-B, 3-D, 4-D, 5- C, 6-C

D. 1-A, 2-B, 3-B, 4-D, 5- C, 6-C

111. Which of these represents true gastro-esophageal (GE) junction tumour?

A. Located between 5 cm and 1 cm above GE junction
B. Located between 2 cm above and 1 cm below GE junction
C. Located between 1 cm above and 2 cm below GE junction
D. Located between 1 cm above and 1 cm below GE junction


5 cm
Type I
Anatomical cardia
1 cm
Type II
–2 cm
Type III
–5 cm

112. Which of the following genetic defect regarding Maturity Onset Diabetes in the Young (MODY) is incorrectly matched?

A. MODY 2 – Glucokinase
B. MODY 3 – HNF-4 alpha
C. MODY 4 – IPF 1
D. MODY 5 – HNF-1 beta

113. 26-year-old woman is being evaluated for a possible inherited disorder. She has a 6-year history of generalized tonic-clonic seizures, and a year ago, she had partial loss of vision due to an occipital infarction. Her mother has chronic intermittent muscle weakness and lactic acidosis, and her maternal uncle has hemiplegia. After further evaluation, all the affected family members are found to suffer from the same inherited disease. Pedigree analysis is shown in the image below. Which of the following is the most likely explanation for the variability of clinical manifestations in the affected family members?

A. Anticipation
B. Heteroplasmy
C. Incomplete penetrance
D. Mosaicism



114. Which of the following statements is true about beta blockers:

1. **Carvedilol is the most widely used beta blocker.**
2. **These are drug of choice in acute decompensated heart failure.**
3. **Pindolol would be less likely to induce severe bradycardia in comparison to the other β-blockers.**
4. **Sotalol can prolong the AP duration leading to torsade's de pointes.**

A. 1, 2, 3, 4

B. 3, 4

C. 1, 3, 4

D. 2, 4

115. Which of the following muscle is supplied by the nucleus present beneath the area marked by the arrow in adjoining image?

A. Risorius
B. Lateral rectus
C. Inferior oblique
D. Medial pterygoid



116. 44-year-old man comes to the emergency department for exertional dyspnea, abdominal distention, and abdominal discomfort. He has a history of alcohol abuse, peptic ulcer disease, and acute pancreatitis. Physical examination shows a distended abdomen with shifting dullness and 1+ ankle edema. A diagnostic paracentesis is performed and shows the following:
Albumin-2.5 g/dL
Neutrophils-200/uL
Blood studies show the following:
Albumin-3.8 g/dL
Glucose-120 mg/dL
Aspartate aminotransferase-50 units/L
Alanine aminotransferase (ALT)-40 units/L
Which of the following is the most likely cause of this patient's ascites?

A. Decreased liver synthetic activity
B. Increased hepatic capillary hydrostatic pressure
C. GI malignancy
D. Pancreatitis

117. 56-year-old man is evaluated for dyspnea, wheezing, and cough productive of yellow sputum. The patient reports that he has been treated with antibiotics when the dyspnea is severe. He has smoked 2 packs of cigarettes daily for the past 25 years. The schematic diagram below is a representation of a bronchial wall. Which of the following ratios is the best pathological index to determine the severity of this patient's disease?

A. A/(A+B+C+D)
B. A/(A+B+C+D+E)
C. C/(B+C+D)
D. C/(A+B+C+D+E)



118. A group of investigators examined the effects of different muscarinic agonists and antagonists on the bladder musculature. The curves presented below were constructed after application of different agents. Curve A represents the effect of bethanechol, a pure cholinergic agonist. Which of the curves most likely represents the effect of bethanechol after application of a reversible competitive antagonist?

A. Curve B

B. Curve C

C. Curve D

D. Curve E



119. 63-year-old man is admitted to the hospital after sufering a cerebral infarction. A swallow study performed during the acute recovery period shows oropharyngeal dysphagia. Despite being maintained on the appropriate dietary restrictions, the patient experiences an episode of vomiting while lying on his back and subsequently develops pneumonia. Which of the following lung regions is most likely to be affected?

A. Anterior segment of the left upper lobe

B. Basilar segments of the right lower lobe

C. Lingular segments of the left lung

D. Posterior segment of the right upper lobe

120. Which of the following anticancer drugs act at the target shown in figure?

A. Erlotinib

B. Ribociclib

C. Dabrafenib

D. Nilotinib



121. A seventeen-year-old patient is brought to the emergency department having just ingested a large overdose of an antihistaminic drug. This drug is a weak base capable of entering most tissues including the brain. Which of the following statements is true:

A. Urinary excretion would be accelerated by administration of NaHCO3, an alkalinizing agent.

B. Urinary excretion would be accelerated by the administration of NH4Cl, an acidifying agent.

C. More of the drug would be ionized at blood pH than at stomach pH.

D. Absorption of the drug would be faster from the stomach than from the small intestine.

122. Identify the incorrect pair of contraceptive dosages

A. Ullipristal for emergency contraception-30mg single dose
B. LNG for emergency contraception 1.5mg two tablets repeated after 12hrs
C. DMPA 150mg im 3 monthly
D. Mirena 52mg LNG with 20ug/d release

123. This is a schematic diagram depicting body structure of which of these helminthes?

A. Loa loa
B. Wuchereria bancrofti
C. Brugia malayi
D. Onchocera volvulus

124. A 12-year-old child is brought to the emergency department after developing blurred vision and confusion. He was playing in the park and ate an unknown fruit. Temperature is 38.9 C (102 F), blood pressure is 110/76 mm Hg, pulse is 120/min, and respirations are 16/min. He is disoriented to time, place, and person. Physical examination shows flushed skin and dry oral mucosa. Both pupils are dilated and nonreactive to light and accommodation. Bowel sounds are decreased. Which of the following drugs can potentially reverse this patient's condition?

A. Atropine
B. Naloxone
C. Physostigmine
D. Thiamine

125. 31-year-old woman is brought to the emergency department by her boyfriend due to confusion. The boyfriend says that the patient "seemed okay" when he left for work this morning, but when he returned in the evening, she was disoriented and "acting like a drunk person"; she had also vomited "all over the bathroom." She has a history of bipolar disorder. Medications include clonazepam, lithium, and quetiapine. Temperature is 37 C (98.6 F), blood pressure is 110/68 mm Hg, pulse is 92/min, and respirations are 16/min. On physical examination, the patient is somnolent and has frequent fasciculations. Pupils are equal and reactive. During evaluation, she has a brief episode of a generalized tonic-clonic seizure. Laboratory results are as follows:
Hemoglobin: 13 g/dL
Leukocytes: 9,000/mm³
Sodium: 136 mEq/L
Potassium: 3.8 mEq/L
Bicarbonate: 28 mEq/L
Creatinine: 1.2 mg/dL
Glucose: 120 mg/dL
Serum lithium 2.8 mEq/L
In addition to supportive measures, which of the following is the best next step in management of this patient?

A. Activated charcoal
B. Calcium gluconate
C. Gastric lavage
D. Hemodialysis

126. A 1-day old baby is evaluated in the emergency department due to poor feeding, repeated bilious vomiting and not passing meconium since birth. Abdominal examination reveals a distended abdomen. Imaging shows the following picture. Identify the condition which is unlikely in this patient?

A. Aganglionosis
B. Anorectal malformation
C. Meconium plug syndrome
D. Meconium ileus

127. Which of the following is likely with ascitic fluid protein 1.9 g/dL and a serum albumin ascites gradient (SAAG) of 1.4?

A. Alcoholic cirrhosis

B. Peritoneal malignancy

C. Constrictive pericarditis

D. Veno-occlusive disease

128. Which of the following drugs reversibly binds the site A?

A. Clopidogrel
B. Prasugrel
C. Ticagrelor
D. Tirofiban



129. In the histology section given, which marked area is involved in relieving pain in response to massage?

A. A
B. B
C. C
D. D



130. A 65-year-old man comes to the emergency department due to severe lower abdominal pain and nausea. He also has not been able to urinate for the past 24 hours. The patient has a history of hypertension and benign prostatic hyperplasia. On examination, a large mass is palpable in the suprapubic area. Multiple attempts at urethral catheterization are unsuccessful, and an ultrasound-guided midline suprapubic cystostomy is planned. Besides the bladder wall, which of the following structures is most likely to be penetrated by the trocar and cannula during the procedure?

A. Anterior abdominal aponeurosis
B. Parietal peritoneum
C. Perineal membrane
D. Visceral peritoneum

Suprapubic abdominal wall



131. A 25-year-old male patient presented with sudden onset breathlessness and a sharp pain in his chest after a road traffic accident. His BP is 85/69mm Hg and HR is 110bpm. The chest X-ray is given below. What is the next step for this patient?

1. **eFAST**
2. **Insert iv cannula and start iv fluids**
3. **Needle thoracostomy**
4. **ICD insertion**
5. **Thoracotomy**
6. **CECT chest**



Options;

A. 1,2,3,4

B. 3,4

C. 2,3,4

D. 1,2,3,4,5,6

132. A female presents to OPD with 6 weeks of amenorrhea complaints of bleeding per vagina and mild abdominal pain. The urine pregnancy test is positive and hCG level is 2800 IU/L. On investigation, mass is seen on the left adnexa measuring 3 x 2 cm. She is hemodynamically stable. How will you manage this patient?

A. Oral methotrexate
B. Single-dose methotrexate injection
C. Serial methotrexate + leucovorin rescue
D. Salpingectomy

133. A 14-year-old girl is brought to the emergency department following a suicide attempt. She swallowed several of her grandfather's theophylline tablets after an intense argument with her parents a few hours ago. The patient reports nausea and headache. Her temperature is 37 C (98.6 F), blood pressure is 114/76 mm Hg, pulse is 88/min, and respirations are 16/min. This patient should be carefully monitored for which of the following?

A. Hypoglycemia and complete heart block
B. Jaundice and elevated liver enzymes
C. Sedation and respiratory depression
D. Seizures and tachyarrhythmias

134. A 19-year-old sex worker was treated 2 days ago for presumed gonorrhea with ceftriaxone and azithromycin. She also has a palpable spleen. Laboratory examination is notable for a Hb of 5 g/dL with normal white blood cells and platelets. Peripheral blood smear shows an excess of spherocytes. Which of the following tests will most likely confirm the diagnosis?

A. ADAMTS-13 activity assay
B. Direct antiglobulin (Coombs) test
C. Flow cytometry
D. G6PD deficiency assay

135. A 47-year-old woman comes to the office due to fever, dysuria, and abdominal pain. The patient has had several urinary tract infections since her teens but has no other medical concerns. Physical examination shows mild suprapubic discomfort. Urinalysis demonstrates pyuria and many bacteria. CT scan of the abdomen is shown in the image below: Which of the following structures has most likely limited the proper ascent of the anomalous organ seen on CT scan?

A. Inferior mesenteric artery
B. Inferior vena cava
C. Persistent urachus
D. Superior mesenteric artery



136. Slit lamp ophthalmological examination of a patient revealed the following. All of the following can cause this except:

A. Quinidine
B. Amiodarone
C. Chloroquine
D. Indomethacin



137. Which among the following species is transmitted through the vector given in the image?

A. R.conorii
B. R.typhi
C. R.akari
D. R.prowazekii

138. A 75-year-old male after a fall in bathroom and the pelvic X-ray is shown below. What will be the position of the left lower limb?

A. Shortened and internally rotated
B. Shortened and abducted
C. Extended, abducted and externally rotated
D. Abducted and internally rotated

139. Area marked A in the given figure contain all the following, except:

A. Zonula occludens
B. Zonula adherens
C. Macula adherens
D. Gap junction



140. 58-year-old, previously healthy man is evaluated due to a 2-month history of early satiety, postprandial epigastric pain and vomiting and unintended weight loss. Physical examination shows cachexia and palpable epigastric mass. Upper gastrointestinal endoscopy reveals a large prepyloric mass causing gastric outlet obstruction; biopsy of the mass is consistent with poorly differentiated adenocarcinoma. Abdominal CT scan reveals thickening of the gastric wall with adjacent fat infiltration, enlarged lymph nodes, and multiple hepatic metastatic lesions. The disease is considered unresectable, and the patient undergoes laparoscopic gastrojejunostomy for palliation of gastric outlet obstruction. Postoperatively, he tolerates tube feeding but experiences worsening weakness. The patient also has frequent episodes of non-sustained ventricular tachycardia. Laboratory results are as follows:

	At admission	Postoperative day 2
Hemoglobin	**9.9 g/dL**	**9.8 g /dL**
Leukocytes	**8,800/mm³**	**10,000/mm³**
Serum chemistry		
Sodium	**130 mEq/L**	**135 mEq/L**
Potassium	**3.7 mEq/L**	**2.8 mEq/L**
Creatinine	**1.1 mg/dL**	**0.8 mg/dL**
Glucose	**90 mg/dL**	**120mg/dl**
Phosphorus	**3.1 mg/dL**	**1.2mg/dl**

Which of the following is the most likely cause of this patient's worsening condition?

A. Hypermetabolic syndrome

B. Nutritional replenishment

C. Paraneoplastic syndrome

D. Sepsis

141. A 40 year old male patient is being evaluated for a suspected brain tumor. A biopsy is obtained whose image has been shown below. Which of the following is the most probable diagnosis based upon the biopsy findings?

A. Pilocytic astrocytoma

B. Glioblastoma multiforme

C. Medulloblastoma

D. Oligodendroglioma



142. A 68-year-old man comes to the OPD for a follow-up appointment. He has a history of advanced melanoma that is unresectable and resistant to adjuvant regimens. The patient recently began receiving monoclonal antibody infusions for advanced melanoma. The monoclonal antibodies block a specific cell surface receptor found on T lymphocytes. As a result, T cells capable of recognizing tumor antigens have improved ability to destroy cancer cells. Which of the following cell surface receptors is most likely blocked by the treatment?

A. CCR5
B. CD4
C. CD19
D. PD-1


Ipilimumab
Atezolizumab
Durvalumab
Avelumab
Cemiplimab
Nivolumab
Pembrolizumab
T cell
MHC II
APC
CD28
B7
CTLA-4
Antigen
MHCI
TCR
Tumor cell
T cell activation
PD-1
PD-L1

143. 52-year-old man comes to the OPD for evaluation of hearing loss in the left ear. The patient works in sales and is frequently on the phone. Over the past several months, he has had difficulty hearing phone conversations in the left ear and must move the headset to the right. He has no pain or vertigo but sometimes feels off-balance when walking at night and holds onto walls. A tuning fork placed on the forehead is lateralised to the the right ear. Air conduction is greater than bone conduction bilaterally. There is decreased sensation to touch on the left side of the face. Which of the following is the most likely cause of this patient's symptoms?

A. Cholesteatoma

B. Herpes zoster oticus

C. C Ménière disease

D. Vestibular schwannoma

144. A 13 year old girl has complaints of recurrent nose bleeds and excessive loss of blood during menstruation. She appears pale. Laboratory studies reveals the following:.
Hemoglobin - 7 g/dL
Platelets - 2 lac/mm3
Bleeding time - Prolonged
PT - Normal
aPTT - Prolonged
Which of the following factors may be deficient in this case?

A. Factor IX

B. Factor XI

C. gp 1b-IX

D. vWF

145. Identify the incorrect statements:
1. Richardson and Koeppe are direct gonioscopes.
2. Narrowing of isopters and the appearance of a blockage within the blind spot are early changes in MS
3.Most common ocular manifestation of COVID-19 is papillary conjunctivitis
4. The Haigis-L regression formula is used for lens power calculation in patients undergoing cataract surgery with a past history of LASIK for myopia.
5. Correct method is to put the eye ointment first, drops 15 min later

A. 1,2,3,4,5
B. 2,3,4
C. 1,3,5
D. 2,3,5

146. Identify the scheme which bears the logo given below

A. Janani Suraksha yojna

B. Janani Shishu Suraksha Karyakram

C. Labour Room Quality Improvement Initiative

D. Vandemataram Scheme

147. Identify the correct pair:

A. Staghorn calculus
B. Putty kidney
C. PUJ obstruction
D. Cortical nephrocalcinosis

A.

B.

C. 

D.

148. The following image depicts an organic poison. Which of the following statements is true about it?

A. Fatal dose is 60-120 seeds
B. Weight of one seed is approximately 10g and was used to measure gold in the past.
C. Symptoms resemble those of cobra bite.
D. Active principle is a toxalbumin.

149. All of the following are true except:

A. Dermis has dense irregular connective tissue
B. Fibrocartilage, bones, tendons have collagen type I
C. Hyaline cartilage has perichondrium with 2–4 chondrocytes in each lacuna
D. Fibres from Inferior olivary nucleus directly reach the purkinje cells in cerebellum, which constitute the largest neurons in human brain

150. You are a medical officer posted in the one stop center of your hospital. You are required to collect evidences from a sexual assault victim. What is the correct order of collection according to the instructions given in the SAFE kit?.1- Cervical swab for culture.2- Vaginal swab.3- Clothing.4- Breast swabs.5- Nail clippings.6- Matted pubic hair cutting.7- Pubic hair combing

A. 3>5>4>6>7>1>2
B. 3>5>4>7>6>2>1
C. 4>5>3>6>7>1>2
D. 3>4>5>76>7>1>2

SAFE kit (sexual assault forensic evidence)

- Step 1- informed consent
- Step 2- history of sexual violence
- Step 3A- outer clothing.● Step 3B- inner clothing
- Step 4A- Debris from hair combing.● Step 4B- Interdigital debris.● Step 4C- Swabs from stain on the body.● Step 4D- Nail scrapings.● Step 4E- Nail clippings.● Step 4F- Scalp hair
- Step 5- Two breast swabs.
- Step 6A- Pubic hair combing.● Step 6B- Cutting of public hair.● Step 6C- Cutting of matted pubic hair.
- Step 7A- Two high vaginal/ cervical swabs.● Step 7B- Two low vaginal swabs.● Step 7C- Two urethral swabs.
- Step 8- Cervical swab for culture.
- Step 9- Vaginal washing.
- Step 10A- Two anal swabs.● Step 10B- Two rectal swabs.
- Step 11- Two oral swabs.
- Step 12- Blood samples for blood grouping, testing for alcohol or intoxication, DNA analysis, testing for HIV, VDRL, HbsAg.
- Step 13- Urine test for pregnancy and drugs.
- Step 14- Medical examination report

151. Identify the correct statements:

1. Miami criteria: 50% reduction in PTH within 10 mins of gland removal

2. Majority of the retrosternal goitres derive their blood supply from mediastinal vessels

3. Purulent peritonitis following a perforated diverticulitis is Hinchey grade 3

4. CECT abdomen in stable patients with eFAST positive is a part of "C" of primary survey

---

A. 1,2,3

B. 1,2,4

C. 1,3

D. 1,3,4

152. A patient has undergone cataract surgery for lens removal and IOL has not been implanted. Instead he is prescribed high power spectacles. Apart from cosmetic blemishes, what other side effects should he not be worried about ?

A. Pin cushion effect
B. Minification of image
C. Roving ring scotoma
D. Jack in the box phenomenon

153. A 7 year old male patient is taken for right side radius-ulna open reduction internal fixation surgery under General anesthesia. As the child seems uncooperative for regional anesthesia, you decide to start general anesthesia induction with an inhalational agent via mask. Within a few minutes the patient starts breath holding and coughing. To find out the cause you check the inhalational agent vaporizer and you immediately turn it off and switch on the non- irritant Inhalational agent. What was the probable Inhalational agent which led to coughing and breath holding?

A. Xenon
B. Desflurane
C. Halothane
D. Sevoflurane

154. A 46 year old male presented to the clinic with a persistent cough, fatigue, weight loss, and night sweats. Xray of the patient is as given below. What is the most likely diagnosis?

A. Streptococcus pneumoniae infection
B. Mycobacterium tuberculosis infection
C. Carcinoma lung
D. Sarcoidosis



155. A 52-year-old man has undergone lung transplantation. Two months after receiving lung transplant, he developed pulmonary symptoms and was diagnosed as having bilateral diffuse interstitial pulmonary pneumonitis and bronchiolitis. Which of the following etiological agent is responsible in this condition?

A. Cytomegalovirus
B. Herpes Simplex virus
C. JC virus
D. BK virus

156. A 3-year-old girl is brought to the office after developing fever and a sore throat. The patient recently entered day care, and similar symptoms have been reported in several of the other children. Physical examination shows exudative pharyngitis and enlarged anterior cervical lymph nodes. A rapid antigen detection test confirms the diagnosis of streptococcal throat infection. Her condition resolves with antibiotic therapy. Several weeks later, she is re-exposed to Streptococcus pyogenes. The bacteria penetrating beyond the surface epithelium are immediately coated with preformed IgG antibodies. Which of the following substances acts in the most similar manner to IgG antibodies to facilitate phagocytosis?

A. Complement C3b

B. Complement C5a

C. Immunoglobulin M

D. Leukotriene B4

157. Following road traffic accident patient suffers polytrauma and is evaluated in the emergency section of the hospital. His pulse rate is 116, respiratory rate is 24, blood pressure of 122/78 mm of Hg and patient is mildly anxious. What is the approximate blood loss patient has following trauma?

A. 750-1500 mL
B. <750mL
C. 1500-2000ml
D. >2000 mL

158. Identify the accessory protein marked in the diagram below

A. Nebulin
B. Alpha actinin
C. Titin
D. Tropomyosin



159. Calculate the corrected reticulocyte count of a patient with haemoglobin of 5 and absolute reticulocyte count of 9%?

A. 4.5%
B. 6%
C. 1%
D. 3%

160. A research laboratory is evaluating a new dipstick test for diagnosing urinary tract infections (UTIs). The new test is compared to urinalysis of a midstream urine specimen (considered the diagnostic gold standard) to establish diagnostic test parameters. The study enrolls 300 patients, of which 100 have a UTI as determined by urinalysis. The new dipstick test is determined to be 70% sensitive and 90% specific for the diagnosis of UTIs. How many false positives are present in the study?

A. 20

B. 30

C. 70

D. 120

161. A 52-year-old female patient present with 4 cm diameter diagnosed breast cancer lesion with ipsilateral axillary and contralateral supraclavicular lymphadenopathy as shown below. As per AJCC system, patient belongs to which stage of breast cancer?

A. T4a N3c M1
B. T4b N1 M1
C. T4b N3c M0
D. T4c N1 M0



162. A woman presented to the clinic with heavy, painful menses that have worsened over the past year. On pelvic examination the uterus is enlarged and tender. MRI scan of the patient's abdomen revealed a finding similar to the image given below. What is the most likely diagnosis?

A. Polyps
B. Leiomyoma
C. Adenomyosis
D. Endometrial hyperplasia

163. A 29 year old female presented with complaints of secondary amenorrhea for the past 6 months. She reports having a regular menstrual cycle in the past and there were no recent changes in weight, exercise, or stress. She does not use any contraceptives and has no significant medical history. On examination, she appears well nourished. Her pregnancy test is negative. What is the next best step in management?

A. Measurement of serum prolactin levels
B. Measurement of serum FSH and LH levels
C. Pelvic USG
D. Endometrial biopsy

164. A patient with metastatic prostate cancer develops hypocalcemia during treatment with a bone antiresorptive agent. Which of the following agents is most likely to result in hypocalcemia?

A. Pamidronate
B. Zoledronic acid
C. Denosumab
D. Ibandronate

165. Which of the following options is incorrectly matched?

1. **Prolapse in pregnancy: Ring Pessary**
2. **Vault prolapse - Sacrospinous ligament suspension**
3. **Cystocele - Colposuspension**
4. **Rectocele - Posterior colporrhaphy**
5. **Prolapse in a female who desires future child bearing-Shirodkar sling surgery**
6. **Prolapse in a 65 year old female with DM and hypertension-Le Fort's colpocleisis**

A. 4,5
B. 3,6
C. 2
D. 3

166. This device shown in the image is used in:

A. Men with suspected urethral stricture
B. Women with recurrent UTI
C. Urine incontinence in pediatric population
D. Suspected VUR

167. A large, multi-country study is conducted to determine the effect of economic development on cancer incidence and mortality. The study uses data obtained from the national cancer registries, along with information regarding per capita gross domestic product as reported by the International Monetary Fund and life expectancy as reported by the World Health Organization. Which of the following best describes the design of this study?

A. Cross-sectional survey

B. Ecological study

C. Nested case-control study

D. Systematic review

168. 23-year-old comes to the ED with the feeling of food being stuck in the throat. Laryngoscopy reveals fish bone lodged in the left piriform recess. During retrieval of the fish bone, a nerve deep to the mucosa overlying the recess is damaged. Which of the following is most likely to be impaired in this patient as a result of his iatrogenic injury?

A. Cough reflex
B. Gag reflex
C. Mastication
D. Salivation

169. Match the following images:

A. Venous ulcer
B. Marjolin ulcer
C. Arterial ulcer
D. Trophic ulcer

1.

2.

3.

4.

A. 1-C, 2-D, 3-B, 4-A
B. 1-D, 2-C, 3-A, 4-B
C. 1-C, 2-D, 3-A, 4-B
D. 1-D, 2-C, 3-B, 4-A

170. The following hematoxylin and eosin-stained biopsy is from which tissue?

A. Tonsils
B. Spleen
C. Lymph node
D. Peyer's patches



171. Identify the correct scenario

1. Loss of sensations over the root of penis after lap hernia repair-Ilioinguinal nerve

2. Hyperesthesia of tongue after parotid surgery-Auriculotemporal nerve

3. Diaphragm palsy after thymoma resection-Phrenic nerve

4. Loss of sensations over the lateral gluteal region after hernia mesh repair-LCNT

A. 1,2,3,4

B. 1,2,3

C. 1,3

D. 2,3,4

172. A therapist observes distinct patterns in her sessions with a 35-year-old woman who has a turbulent history with relationships. Initially, the woman sings praises about the therapist, declaring her as the best thing that's happened to her. However, after a few sessions, she arrives agitated, accusing the therapist of not genuinely wanting to help her. These extreme shifts in perception occur frequently, creating a turbulent therapeutic environment. Which defense mechanism is predominantly being exhibited by this patient?

A. Reaction formation
B. Denial
C. Projection
D. Splitting

173. The anesthetist feels a snap while performing an epidural catheter insertion. This is due to piercing of which structure?

A. Ligamentum flavum
B. Supraspinous ligament
C. Dura mater
D. Posterior longitudinal ligament

174. A 32-year-old female presents to the dermatology clinic concerned about white patches on her skin that she noticed since childhood. Physical examination reveals depigmented patches with a symmetrical distribution, especially prominent on the ventral aspect of her body and mid-forehead. White forelock is noted. She mentions that her grandfather had similar patches since his birth. She also reports that a cousin on her father's side has been recently diagnosed with rheumatoid arthritis. Given the patient's history and clinical presentation, which of the following describes the underlying cause of her condition?

A. A polygenic disorder with progressive depigmentation.
B. An autosomal dominant disorder involving the C-kit protooncogene.
C. An autoimmune condition with progressive depigmentation.
D. A sporadic condition without a genetic link

175. A 73-year-old man comes to the emergency department with unstable angina. He undergoes coronary angiography via the femoral approach. A stent is placed in the right coronary artery, and the patient is discharged. He returns to the emergency department 2 days later with blue discoloration of his right toe. He has pain and mild tingling in the affected toe. On physical examination, the right toe appears cyanotic, and there is livedo reticularis affecting the right thigh. Serum creatinine is 2.8 mg/dL (preoperatively it was 1.0 mg/dL). Which of the following histopathologic findings would most likely be seen on biopsy of this patient's kidney?

A. Cholesterol clefts in arterial lumen

B. Glomerular crescent formation

C. Extensive necrosis of proximal tubular cells

D. Tubular obstruction with rate crystals

176. A 60 year old male with history of long standing goitre visits the OPD. Surgeon performs this manoeuver- which makes the patient breathless. This finding implies?

A. Goitre is retrosternal
B. Malignant transformation
C. Post operative airway risk
D. Infiltration of carotid artery

177. In which of the following study designs, the Outcome of interest has already taken place when the study is started?

A. Prospective cohort study
B. Retrospective cohort study
C. Nested case control study
D. Both (b) and (c)

178. A surveillance study is conducted to assess the long-term efficacy and safety of a drug currently being used to treat patients with heart failure. Researchers enroll 8,300 patients with heart failure. The patients receive the drug once daily for 6 months. The results show significant clinical improvement, but severe hypernatremia is observed in 23 patients. The study publication recommends a lower dose of the drug in patients with baseline normonatremia and hypokalemia to prevent hypernatremia. Which of the following best characterizes this type of study?

A. Phase I clinical trial
B. Phase II clinical trial
C. Phase III clinical trial
D. Phase IV clinical trial

179. A patient presents to the emergency room with symptoms of fever, increased respiratory rate, elevated heart rate, and low blood pressure. The physician suspects the patient may have SIRS. Which of the following is not a criteria for SIRS?

A. Systolic blood pressure <90 mmHg
B. Temperature >38 degree Celsius or <36 degree Celsius
C. Respiratory rate >20 bpm
D. Heart rate >90 min

180. A 23-year-old man with HL was treated successfully with the ABVD regimen. Which of the following drugs used in the treatment of this patient is cell cycle specific ?

A. Alkylating agents
B. Corticosteroids
C. Antimetabolites
D. Plant alkaloids


Bleomycin
Microtubule inhibitors
Taxanes
Vinca alkaloids
Topoisomerase inhibitors
Etoposide
Teniposide
Irinotecan
Topotecan
Antimetabolites
Cladribine*
Cytarabine
5-flurouracil
Hydroxyurea
Methotrexate
Pentostatin
Thiopurines
*Cell cycle
Nonspecific
Cell cycle-independent drugs
Platinum compounds
Alkylating agents:
Anthracyclines
Busulfan
Dactinomycin
Nitrogen mustards
Nitrosoureas
Procarbazine
XX
G1
Double check repair
Mitosis
M
Cytokinesis
Interphase
DNA synthesis
Duplicate cellular content
S
G1
G0
Resting
Rb, p53 modulate
G restriction point

181. A patient is induced with Thiopentone IV through one of the antecubital veins. The patient complains of severe pain engulfing his whole hand. What is the next line of management?

A. Give IV propofol through same needle
B. Insert another iv line immediately
C. IV lignocaine through same needle
D. Remove the needle immediately

182. A 60-year-old male is brought to the emergency department by ambulance after a road traffic accident. He doesn't know where he is, he is unable to identify his relatives and is in a state of confusion. On examination, he has eye-opening only on painful stimuli and can localize the site of painful stimuli. What is the GCS score of this patient?

A. E2V3M5

B. E2V4M5

C. E3V3M6

D. E3V4M6

183. A 48-year-old male presented to the clinic with complaints of acute abdominal pain, nausea, and vomiting. The pain is sudden onset, and severe, in the epigastric and umbilical regions, referred to the back and relieved on bending forwards. He has a past history of chronic alcoholism. On examination, the abdomen is tender with guarding and rigidity, rebound tenderness is present and mild jaundice is evident. Which of the following is the best investigation for this patient?

A. Liver function test

B. Serum amylase

C. USG abdomen

D. CECT scan

184. A 32-year-old female presented to the clinic with complaints of heart racing, diarrhea and intolerance to hot weather. She has a known history of iodine deficiency goiter and is taking iodine supplementations. She has no history of flu-like illness in the past weeks. Her blood results on the recent follow-up are as follows: high T3, high T4 and low TSH. What is the most likely diagnosis?

A. De Quervain thyroiditis

B. Wolf-Chaikoff effect

C. Jod-Basedow effect

D. Hypothyroidism

185. Identify the marked layer:

A. Outer plexiform layer
B. Inner plexiform layer
C. External limiting membrane
D. Internal limiting membrane



186. Identify the incorrect pair

A.

B.

C.

D.

A. C4c

B. C1

C. C4a

D. C4b

187. A group of sports physicians plans to conduct a case-control study to investigate a possible association between adolescent idiopathic scoliosis (AIS) and sacroiliac joint (SIJ) dysfunction in young athletes. The case group will consist of young athletes who were diagnosed with AIS during a regular checkup by a sports physician. Which of the following is the most appropriate control group for this study?

A. Young athletes with a diagnosis of AlS and SIJ
B. Young athletes with a diagnosis of AlS but not of SIJ
C. Young athletes with a diagnosis of AlS irrespective of SIJ status
D. Young athletes with no diagnosis of AIS irrespective of SIJ status

188. Identify the correct statements:

1. **Thelarche is the first sign of puberty in females.**
2. **Maximum growth occurs in Tanner stages 4 and 5 in males**
3. **Penile enlargement is the first sign of puberty in males**
4. **Peak growth velocity always precedes menarche in females**

A. 1,2,4

B. 1,2,3,4

C. 2,3,4

D. 1,2,3

Match the following images with the correct name of the manouever

A. Pawlick grip
B. Pelvic grip
C. Rubin manoeuvre
D. McRobert manoeuvre
E. Ritgen manoeuvre
F. Burns Marshall manoeuvre
G. Mauriceau–Smellie–Veit maneuver



1

2

a) 1-D, 2-B, 3-G, 4-F
b) 1-D,2-A, 3-E, 4-F
c) 1-C,2-A, 3-G, 4-C
d) 1-G,2-B, 3-C, 4-G



3



4

190. Correctly identify the type of suturing shown below.

A. Subcuticular sutures
B. Interrupted mattress sutures
C. Interrupted simple sutures
D. Continuous sutures



191. Identify the correct statements:

1. **Social smile – 2months**
2. **Walk without support -12months**
3. **Babbling gibberish-15months**
4. **Walk upstairs with alternate foot – 4yrs**
5. **Copy a circle-2yrs**

A. 1,2,3,4

B. 1,3,4,5

C. 1,3

D. 1,3,5

192. A 65-year-old male with a history of atrial fibrillation and recent coronary artery bypass graft surgery presents with bleeding from the surgical wound site. Laboratory tests reveal a normal platelet count and coagulation panel. Thromboelastography (TEG) is ordered to assess the patient's clotting profile. Which of the following TEG tracings is most consistent with the patient's presentation?

A. Reduced R-time, increased angle, increased MA, increased LY30

B. Prolonged R-time, decreased angle, decreased MA, decreased LY30

C. Prolonged R-time, increased angle, decreased MA, increased LY30

D. Reduced R-time, decreased angle, increased MA, decreased LY30

TEG	ROTEM	Coagulopathy	Treatment options
R 10–14 min	ExTEM CT 80–100 s InTEM CT 200–240 s	Coagulation factors ↓	FFP 20 ml/kg
R >14 min	ExTEM CT >100 s InTEM CT >240 s	Coagulation factors ↓↓	FFP 30 ml/kg
FF_{MA} 7–14 mm	FibTEM MCF 6–9 mm	Fibrinogen ↓	FFP 20 ml/kg or cryoprecipitate 3 ml/kg or fibrinogen concentrate 20 mg/kg
FF_{MA} 0–7 mm	FibTEM MCF 0–6 mm	Fibrinogen ↓↓	FFP 30 ml/kg or cryoprecipitate 5 ml/kg or fibrinogen concentrate 30 mg/kg
MA 45–49 mm and FF_{MA} >14 mm	ExTEM A_{10} 35–42 mm and FibTEM ≥10 mm ExTEM MCF <50 mm and FibTEM ≥10 mm	Platelets ↓	Platelets 5 ml/kg
MA <45 mm and FF_{MA} >14 mm	ExTEM A_{10} <35 mm and FibTEM ≥10 mm	Platelets ↓↓	Platelets 10 ml/kg
Ly30 >8%	ExTEM Li30 <94%	Hyperfibrinolysis	TXA 1–2 g IV or 10–20 mg/kg
Difference in R Hep TEG vs. standard TEG R >2 min	InTEM CT/HepTEM CT >1.25	Heparinazation	Protamine 50–100 mg or FFP 10–20 ml/kg

FFP, fresh frozen plasma; S, second; TXA, tranexamic acid.
Thrombelastograph (TEG): R, reaction time (normal range 3–8 min); MA, maximum amplitude (normal range 51–69 mm); Ly30, hyperfibrinolysis after 30 min (normal range <8%; FF_{MA}, functional fibrinogen MA (normal range 14–24 mm).
Thrombelastometry (ROTEM): CT, clotting time (normal range CT_{EXTEM} 38–79 s, CT_{INTEM} 100–240 s, CT_{HEPTEM} 100–240 s); A10, amplitude after 10 min (normal range $A10_{EXTEM}$ 43–65 mm, $A10_{FIBTEM}$ 7–23 mm); MCF, maximum clot firmness (normal range MCF_{EXTEM} 50–72 mm); Li30, hyperfibrinolysis after 30 min (normal range LI30 94–100%).

193. A 34-year-old man presents to the emergency department with severe abdominal pain and distension. On examination, his vital signs reveal a heart rate of 120 beats per minute and blood pressure of 80/50 mmHg. Abdominal examination reveals diffuse tenderness and distension. The plain abdominal radiograph is shown below. What is the most likely diagnosis?



A. Crohn's disease

B. Ulcerative colitis

C. Colorectal carcinoma

D. Toxic megacolon

194. A Female had a thorn prick 5 years ago. She presented with a slowly growing 2 × 2 cm verrucous lesion which on KOH mount showed the following image. Identify the condition?

A. Phaeohyphomycosis
B. Blastomycosis
C. Sporotrichosis
D. Chromoblastomycosis

1.



195. A 13-year-old girl is brought to the office for evaluation of acne. She began to develop acne over her forehead and chin 4 months ago. She used over-the-counter topical salicylic acid, but the acne did not improve. Now, the patient has painful acne over her face and upper chest. She has not had a menstrual period. The girl has grown 10.1 cm in the last year; her height and weight are at the 70th percentile for age. Vital signs are normal. Nodulocystic acne is present across the face and upper chest. There is no breast bud development. The abdomen is nontender and has no palpable masses. External pelvic examination shows the clitoris protruding from the clitoral hood and bilateral masses in the labia majora. Which of the following is the most likely diagnosis in this patient?

A. 5-alpha-reductase deficiency

B. Androgen insensitivity-syndrome

C. Nonclassical congenital adrenal hyperplasia

D. PCOD

196. A new antiarrhythmic drug is found to be effective against both atrial and ventricular arrhythmias. Its effect on action potential is shown in the Figure. The effect of this new drug is most similar to?

A. Lignocaine
B. Propanolol
C. Encainide
D. Quinidine



197. A 56-year-old man presented with the sudden onset of a severe headache. He was awake, but feeling uncomfortable because of headache. His initial plain CT head was normal. Lumbar puncture was performed and CSF was shown below. What is the next step?

A. Repeat plain CT head
B. Digital Subtraction Angiography
C. Ceftriaxone + Vancomycin
D. Aspirin

198. A 24-year-old man is brought to the emergency department due to weakness, lethargy, nausea, and dizziness. He took fifty 325-mg acetaminophen tablets approximately 6 hours ago. Vital signs are normal. Pupils are equal, round, and reactive to light; there is mild epigastric tenderness. Laboratory results include an elevated serum acetaminophen level. Intravenous infusion of N-acetylcysteine is started. This treatment is most likely to improve the patient's condition by which of the following mechanisms?

A. Activating hepatic glucuronidation enzymes

B. Competitively blocking drug receptor sites

C. Increasing the amount of intrahepatic glutathione

D. Increasing renal excretion of drug molecules

199. Identify the true statements about normal fetal development:

1. Endovascular type of extravillious trophoblasts invade the spiral arteries lumen

2. Blood in inferior vena cava has lower oxygen concentration compared to superior Vena cava

3. Functional closure of the fetal circulatory adjustments occurs in the following sequence: Ductus venosus > Foramen ovale > Ductus arteriosus

4. The main factor for the ductal closure postnatally is an increase in partial pressure of oxygen (paO2)

A. 1,2,3,4
B. 1,2,3
C. 1,3,4
D. 2,3,4

200. What does IP in the figure shown stands for?

A. Drug is an instant release preparation
B. Meant for intra pulmonary use
C. The standards according to Indian Pharmacopoeia
D. Drug tested preclinically via Intraperitoneal route

Thank You

BTR Mega Test and Discussion

Dr. Zainab Vora

Cerebellum
Get the balance right

1. A 54-year-old woman comes to the clinic due to difficulty hearing for the past few weeks. During the neurologic examination, the physician assesses her hearing using a vibrating tuning fork. The handle of the tuning fork is placed on her left mastoid process until the sound is no longer audible. The tines are then quickly placed near the patient's left auditory meatus, and she reports hearing no sound. When the handle of the vibrating fork is placed on the middle of her forehead, she hears the vibration more strongly in her left ear than her right. This patient is most likely experiencing which of the following types of hearing loss?

A. Conductive loss in left ear
B. Conductive loss in right ear
C. Sensorineural loss in both ears
D. Sensorineural loss in left ear



2. Which of the following method of disinfection and sterilization can also kill spores?

1. Glutaraldehyde
2. Ethylene oxide
3. Pasteurization
4. Orthophthalaldehyde

A. 1, 3 and 4

B. 1, 2 and 4

C. 2,3 and 4

D. 1,2,3 and 4

Cerebellum
Get the balance right

Disinfectant	Activity	Advantages	Disadvantages	Recommendation
Glutaraldehyde	Broad spectrum microbicidal and sporicidal	Good compatibility	Requires activation Produces irritant fumes	For fibroscope and respiratory equipment
Orthophthaldehyde	Broad spectrum microbicidal and sporicidal	No activation required No fumes	Costly Stains equipment	For scopes Alternative to glutaraldehyde
Iodine compounds	Microbicidal spares M Tuberculosis and spores	Rapid action	Corrosive to metals, plastic and rubber Stains items	As an antiseptic
Alcohols	Wide microbicidal activity Non sporicidal	Non staining	Flammable	Hand disinfection For endoscopes
Phenols	Wide microbicidal activity Non sporicidal	Easily available Low cost	Irritant to skin Depigmentation	As surface disinfectant
Quaternary Ammonium compounds	Microbicidal spares M Tuberculosis, Not sporicidal, Not virucidal	Less irritant Good detergent property	Occupational asthma	As surface disinfectant For non-critical items
Peracetic acid	Broad spectrum microbicidal and sporicidal	No activation required Wide compatibility	Expensive Irritant to eye and skin	For fibroscopes
Chlorines	Wide microbicidal activity Non sporicidal	Low cost Fast acting	Corrosive to metals	Surface disinfectant To clean blood and body fluid spills
Hydrogen Peroxide	Broad spectrum microbicidal and sporicidal	No activation required	Serious eye damage Incompatible with some metals	Fogging of operating room For endoscopes
Formaldehyde	High level disinfectant	Non corrosive	Pungent odor and irritant fumes Carcinogenic	Withdrawn from use

3. Which of the following statements regarding reducing activation energy are true:

1. ΔGf is lowered
2. Stability of transition state is reduced
3. It depends upon complementarity of active site of enzyme and substrate
4. Conformational change may occur at the active site of enzyme at the time of binding to substrate

Cerebellum
Get the balance right

A. 1,2,3,4
B. 2,3,4
C. 3,4
D. 1,4



activation energy ↓

enzyme

Gibbs free energy constant

4. A 41-year-old woman is being evaluated for peripheral edema. The patient has gained 6.8 kg over the past 8 weeks. Her urine is "frothy." Blood pressure is 140/90 mm Hg and pulse is 80/min. Examination shows generalized edema. Heart sounds are normal. The abdomen is soft and nontender. Serum creatinine is 1.1 mg/dL. Urinalysis shows 4+ protein, 0-3 leukocytes/hpf, and oval fat bodies. Serum contains IgG4 antibodies to the phospholipase A2 receptor (PLA2R). Which of the following is the most likely diagnosis?

A. Focal segmental glomerulosclerosis ← Finnish type nephrin 50%.
B. Membranous nephropathy ✓
C. Minimal change disease ✗
D. Mixed cryoglobulinemia ✗

5. A 40-year-old nulligravida woman comes to the office with a 2-month history of worsening right pelvic pain. She experiences the pain daily, and its intensity does not vary during her menstrual cycle. In addition, the patient's waist size has increased despite a decreased appetite. Pelvic examination shows an irregularly shaped, fixed adnexal mass. One week later, she undergoes surgery to remove a large ovarian neoplasm. Within the true pelvis, the surgeon can most likely palpate the right ureter immediately anterior to which of the following structures?

A. Gonadal vein
B. Inferior vena cava
C. Internal iliac artery
D. Uterine artery ← ureter lies post to uterine A

QQ



6. A 45-year-old male patient presents to their primary care physician with a complaint of dysphagia. He reports that he experiences a sensation of food getting stuck in his throat when he eats. He also reports some mild chest discomfort with exertion. The patient has a history of Down syndrome and congenital heart disease. Which imaging modality is typically used to diagnose dysphagia lusoria?

→ Ba swallow

ARSA - post to esophagus

A. Chest X-ray
B. Electrocardiogram
C. Echocardiogram
D. Computed tomography angiography

Cerebellum
Get the balance right

7. A 35-year-old male, diagnosed with squint presented to the ophthalmology department for regular check-up. On examination, marked muscle was found affected. At which level does the nerve supply for the marked structure arise?

A. Red nucleus ← sup midbrain

B. Sub thalamic nuclei

C. Decussation of pyramidal tract

D. Olivary nucleus

MR → 3rd CN



Cerebellum
Get the balance right

8. A 10-year-old boy came with complaints of anaemia, weakness, with low weight to height ratio for his age profile. On examination, urine sample shows elevated level of orotic acid. Further study reveals low uridine monophosphate (UMP) synthase activity. What can be the best treatment to administer?



A. Adenine
B. Uridine
C. Hypoxanthine
D. Thymidine

Cerebellum
Get the balance right


PRPP
Glycine
De Novo
biosynthesis
AMP deaminase
GMP reductase
ADSL
AMPS
ADSS
IMP
IMPDH
XMP
GMPS
AMP
GMP
Inosine
Salvage synthesis
Hypoxanthine

9. A patient presented with fever, dyspnea, palpitations, and tachycardia. The echocardiography revealed vegetation in the septal leaflets. Tissue sections from them reveal the following findings as shown in the image. What is the most likely implicated agent?

Q2



A. Pigmented septate hyphae at 90 degrees, Mucormycosis

B. Right angle branching septate hyphae, Fusarium

C. Acute angle branching hyphae, Rhizopus

D. Acute angle branching septate hyphae, Aspergillus

A sp

10. 15-year-old boy comes to the OPD due to right ear itching and discomfort for the past several days. He has no fever or hearing loss but has noted scant drainage of thin, whitish fluid. The patient has been taking swimming lessons at a local gym. On examination, there is no redness around the ear, but gentle traction of the pinna elicits pain. During inspection of the external auditory canal, a speculum is inserted into the meatus in close contact with its posterior wall, causing the patient to suddenly become lightheaded and faint. He recovers spontaneously within a few minutes with no residual confusion. Which of the following nerves was most likely irritated during the procedure?

A. Facial nerve

B. Jacobson nerve

C. Arnold nerve

D. Auriculotemporal nerve



11. Identify the Diagram shown in the image below?



A. PERT management technique

B. Pie chart model

C. Epidemiological triad model

D. Sufficient component cause model



12. A 45-year-old woman is brought to the hospital after she collapsed during an airshow on a hot summer day. She is receiving amitriptyline for fibromyalgia. Temperature is 40.5 C (104.9 F), blood pressure is 90/60 mm Hg, pulse is 110/min, and respirations are 22/min. The skin is warm and red. She is disoriented. Neurologic examination shows no focal findings. If this patient's medication contributed to her current condition, which of the following mechanisms is most likely responsible?

A. Impaired cardiovascular response to heat stress

B. Impaired dissipation of body heat

C. Increased heat production by increasing muscle tone

D. Increased peripheral vasoconstriction



13. Identify the correct statement

1. Type I collagen is seen in bone and elastic cartilage
2. Reticular fibres are present in lymph nodes, bone marrow and ~~thymus~~
3. Right-left movement is by atlanto-axial joint
4. Costo-transverse joints are plane synovial joints

A. 1,2,3

B. 2,3,4

C. 3,4

D. 1,2,3

Cerebellum
Get the balance right

14. Identify the false statement from the following:

A. True positive fistula test indicates perilymph fistula (T)
B. False positive fistula test indicates congenital syphilis
C. True negative fistula test indicates late stages of Meniere's disease → False (+)
D. False negative fistula test indicates dead labyrinth / cholesteatoma covering fistula



15. A 4-year-old girl is brought to the clinic due to an inward deviated right eye. Examination shows a pale red reflex in the right eye relative to the left eye. Dilated funduscopic examination reveals a well-circumscribed white mass within the retina. Genetic analysis shows a germline mutation in the patient's cells leading to development of the mass. Which of the following is the most likely function of the protein affected by this patient's gene mutation?

A. Activation of receptor tyrosine kinase

B. Conversion of GTP to GDP

C. Opposition of p53 activity

D. Prevention of the G1/S cell cycle transition



Cerebellum
Get the balance right

16. Identify the true statements:

1. At physiological pH, arginine has a positive charge
2. Selenocysteine is coded by UGA
3. Heme synthesis occurs in BM, RBCS and hepatocytes
4. Vitamin C cannot be produced in humans due to lack of L-gulonolactone oxidase

A. 1,2,3,4

B. 3,4

C. 1,2,4

D. 2,3

Cerebellum
Get the balance right

17. All of the following muscles have parallel oriented fibers except:

A. Sartorius
B. Rectus abdominis
C. Sternohyoid
D. Tibialis anterior

Cerebellum
Get the balance right

Muscles with parallel fasciculi

- Muscles in which fasciculi are parallel to the line of pull and have greater degree of movement.
- They are further classified based upon the shape:
  - *Quadrilateral*: Thyrohyoid
  - *Strap-like*: Sternohyoid and sartorius
  - *Strap-like with tendinous intersections*: Rectus abdominis
  - *Fusiform*: Biceps brachii, digastric

Muscles with oblique fasciculi

- Muscles fasciculi are arranged oblique to the line of pull. This increases the power of the muscle, but reduces the range of movement.
- They are further classified into:
  - *Triangular*: Temporalis, adductor longus, trapezius
  - *Unipennate*: Flexor pollicis longus, extensor digitorum longus
  - *Bipennate*: Rectus femoris, flexor hallucis longus
  - *Multipennate*: Tibialis anterior, subscapularis, deltoid (acromian fibers).

18. A 25-year-old homeless man with a history of schizophrenia is brought to the hospital by local police. He destroyed a television set at a local electronics store and then became extremely agitated and violent when employees attempted to intervene The patient's speech is difficult to follow, and he is distracted, seemingly listening to voices that only he can hear. He is admitted to the psychiatric ward and given haloperidol, which calms him down. Later that evening, however, he walks to the nurses' station and says, "What's happening to me?" The patient is upset and refuses to sit down, pointing to his neck. Physical examination shows a sustained contraction of his neck to the right side. Which of the following is the most appropriate next step in management?

Acute dystonia

A. Haloperidol ✗

B. Diphenhydramine ← anticholinergic DOC

C. Levodopa ✗

D. Lorazepam ✗✗

Cerebellum
Get the balance right

19. A 35-year-old, previously healthy man is evaluated immediately after accidental exposure to ionizing radiation at a fluoroscopy manufacturing facility. The patient is currently asymptomatic. Vital signs are normal. Physical examination shows no abnormalities. If radiation exposure is significant, which of the following cells are most likely to be affected first?

A. Bladder epithelial cells

B. Cardiac myocytes

C. Intestinal crypt cells

D. Renal proximal tubular cells

20. Match the following:

1. Stimulation of this area results in turning the eyes and head to the contralateral side
2. A lesion here results in nonfluent, telegraphic speech
3. Ablation in this area results in a contralateral upper homonymous quadrantanopia
4. A lesion here results in fluent speech with neologisms
5. A lesion of this area is characterized by finger agnosia, dyscalculia, dysgraphia, and dyslexia
6. Destruction of this area results in an aphasia characterized by fluent speech, good comprehension, and poor repetition
7. Lesions in this gyrus result in contralateral astereognosis
8. Lesions in this gyrus result in contralateral spasticity

Cerebellum
Get the balance right

A. 1-H, 2-F, 3-G, 4-D, 5-E, 6-C ,7-A, 8-B

B. 1-H, 2-G, 3-F, 4-E, 5-D, 6-C ,7-B, 8-A

C. 1-H, 2-G, 3-E, 4-F, 5-D, 6-B ,7-C, 8-A

D. 1-D, 2-G, 3-F, 4-E, 5-H, 6-C ,7-B, 8-A



21. A 5-month-old girl was brought to the casualty with seizures. The paediatrician tries to do CSF sampling. What are the structures punctured by the paediatrician while piercing through anterior fontanelle?

A. Scalp, dura, arachnoid

B. Scalp, epicranium, endocranium and dura

C. Scalp, synchondral membrane, dura, arachnoid

D. Pericranium, dura, arachnoid

22. A research group conducted a study to compare the levels of creatine kinase-MB (CK-MB) between type 2 diabetes mellitus (T2DM) patients given statin therapy. Participants were divided into 4 groups based on treatment. Groups I, I, and III consisted of T2DM patients who had been given statin therapy (atorvastatin, simvastatin, rosuvastatin, respectively) for at least 6 months. Group IV consisted of T2DM patients who had not been given statin therapy. Which of the following statistical tests is most appropriate to compare the CK-MB levels between Groups I, I, III, and IV in this study?

A. Analysis of variance (ANOVA)

B. Chi-square test

C. Unpaired t-test ✗

D. Paired t-test ✗

Cerebellum
Get the balance right

23. Identify the condition showing the following variation in plasma proteins:

A. Multiple myeloma
B. Hepatic cirrhosis
C. Nephrotic syndrome
D. TB



24. Identify the incorrect pair:

1. Talocalcaneonavicular joint – Saddle joint
2. Malleo-incudal joint-Ball and socket joint
3. Atlanto-axial joint-Pivot joint
4. Acromio-clavicular joint-Plane joint

A. 1,3,4

B. 2,3,4

C. 1,2

D. 2,4

Cerebellum
Get the balance right

25. Identify the organism from the life cycle

A. Ancylostoma duodenale

B. Strongyloides stercoralis

C. Ascaris lumbricoides

D. Enterobius vermicularis



26. A 50-year-old man is scheduled to undergo a colonoscopy for routine colorectal cancer screening. He has no abdominal pain, gross blood in the stool, constipation, change in stool caliber, or unintentional weight loss. He also has no family history of colon cancer. The procedure is uncomplicated, but the patient is found to have a small mucosal lesion (~1 cm) in his sigmoid colon. The lesion is biopsied and its histology is shown in the image below. The pathogenesis of this patient's lesion most likely began with a mutation involving which of the following genes?

A. APC

B. STK11 ← Peutz-Jeghers

C. SMAD4 ← Juvenile polyp

D. MSH2 ← HNPCC

FAP ⇒ > 100 polyps

villions Adenome



27. A 24-year-old male is brought to the psychiatry emergency department by his family members. They report that he has been muttering to himself, exhibiting suspicious behaviour, and believing that neighbours are conspiring against him by video recording him. He has not attended work for the last three months. Over the past month, he has become more withdrawn, speaking minimally, and has been observed standing in his room at the same place for hours. In the last two days, the patient has stopped speaking completely and has refused food and water. In the emergency department, the patient did not respond to any questions and stood as given in the image below for 30 minutes. What is the best next step in the management of this patient?

A. Risperidone
B. IV lorazepam
C. Electroconvulsive therapy (ECT)
D. Haloperidol

Cerebellum
Get the balance right

28. Which of the following is NOT one of the three clinical diagnostic criteria for bacterial vaginosis described by Amsel?

A. Abnormally high vaginal pH

B. Presence of an abnormal discharge and erythema of the vagina

C. Clue cells seen on vaginal saline preparation by light microscopy

D. Characteristic fishy odour release with addition of potassium hydroxide to vaginal secretions

Cerebellum
Get the balance right

29. Identify the incorrect pair

#CC

A. Montenegro test – Kala azar
B. Frenkel test – ~~Malar~~ia → Toxoplasma
C. Fairley test – Schistosomiasis
D. Bachman test – Trichinellosis

Cerebellum
Get the balance right

30. Match the following

1. Main elevator of mandible
2. Superior head originates from infratemporal surface and infratemporal crest of the greater wing
3. Receives supply from main trunk of trigeminal nerve
4. Originates from zygomatic arch and inserts on the angle of mandible

A. A-1, B-2, C-3, D-4

B. A-1, B-4, C-2, D-3

C. A-2, B-1, C-3, D-4

D. A-1, B-4, C-3, D-2





31. 35-year-old man with a history of bipolar and substance use disorders comes to the emergency department due to depression, auditory hallucinations, and suicidal ideation. His medications include lithium and escitalopram. The patient has a history of 5 psychiatric hospitalizations and 2 past suicide attempts, including overdose on his medications and attempted hanging. The patient has been using "anything I can get my hands on" because his depression is unbearable. He is hospitalized and placed on suicide precautions. His dose of escitalopram is increased to target his depression, and risperidone is added to treat the hallucinations. His lithium level is 1.0 mEq/L. On the second day of hospitalization, the patient reports muscle pains, abdominal cramping, nausea, and diarrhea. His temperature is 37.2 C (99 F), blood pressure is 130/85 mm Hg, and pulse is 84/min. The patient is alert and restless, and his pupils are dilated bilaterally. Bowel sounds are hyperactive and neurologic examination is normal. Which of the following is the most likely explanation for his symptoms?

A. Serotonin syndrome ← (+) CLONUS
B. Cocaine withdrawal ✗
C. Lithium toxicity ✗
D. Opioid withdrawal ✓

32. Identify the true statements:

1. IVC passes through the central tendon of diaphragm
2. Left superior intercostal vein drains into the azygos vein
3. There are two dorsal interossei at the middle finger
4. Superior relation of epiploic foramina is formed by the hepatoduodenal ligament and right adrenal

A. 1,2,3,4

B. 1,2,3

C. 1,3

D. 2,4

Cerebellum
Get the balance right

33. Identify the false statements:

1) Rickettsial morula in a monocyte is diagnostic of Ehrlichia chaffeensis

2) E6, E7 are targets of vaccine against HPV

3) SARS CO-2 specimen handling for molecular testing needs BSL-2

4) Peptostreptococcus causes scromboid fish poisoning

A. 2, 4

B. 1,2,4

C. 3,4

D. 2,3

Cerebellum
Get the balance right

34. A 45-year-old previously healthy man comes to the office for evaluation of a neck lump. The patient first noticed the swelling a month ago while shaving, and it has progressively enlarged. On physical examination, there is a nontender, firm nodule in the upper portion of the left thyroid lobe. Laboratory studies reveal a normal serum TSH and elevated serum calcitonin levels. Neck ultrasonography shows a 2-cm hypoechoic thyroid nodule with microcalcifications. Fine-needle aspiration of the nodule reveals spindle-shaped cells in an amorphous background. Which of the following gene alterations most likely contributed to this patient's current condition?

MTC

A. Activation of ALK

B. Activation of RET

C. Inactivation of MEN1

D. Activation of GNAS

Cerebellum
Get the balance right

35. Identify the distribution represented by ‘diagram 3’ in the Image below?

A. Normal distribution
B. Normal distribution with outliers
C. Positive skew
D. Negative skew



Cerebellum
Get the balance right

36. If changes made in some of the sequence of amino acids in the chain of human proinsulin, which of the following changes will not make any difference in the biochemical activity of the molecule?

A. Breaking disulfide linkage
B. Interchange of A1 and A4
C. Interchange of B29 and B30
D. Interchange of A5 and A6



37. Which procedure is shown in the following picture?



A. Infratrochlear block
B. Mandibular nerve block
C. Anterior ethmoidal block
D. Maxillary nerve block

Cerebellum
Get the balance right


nfratrochlear nerve
Supratrochlear nerve
External branch of anterior ethmoidal nerve
Infraorbital nerve

38. Match the following with the appropriate diagnosis:

A. Lichen planus
B. Anti-synthetase syndrome → anti JO 1
C. Osteoarthritis
D. Dermatomyositis

A. 1-a, 2-d, 3-c, 4-b
B. 1-a, 2-d, 3-b, 4-c
C. 1-d, 2-a, 3-c, 4-b
D. 1-d, 2-a, 3-c, 4-d

39. The nerve being tested here is injured most likely at what site?

A. Between two heads of pronator teres
B. At the level of supinator muscle
C. Radial groove
D. Medial epicondyle



Cerebellum
Get the balance right

40. Mr. Ramesh, a 55-year-old male patient, has been diagnosed with laryngeal malignancy and is scheduled to undergo modified radical neck dissection (MRND). The surgery is being performed in a sitting position to provide the surgeon with better access to the operative field. The anaesthetic team has taken all necessary precautions to prevent any complications during the surgery. During the surgery, while ligating the internal jugular vein, the patient's end-tidal carbon dioxide (etCO2) dropped from 40 mmHg to 14 mmHg. The patient suddenly developed hypotension along with cardiac arrhythmia, and the surgical team immediately recognized the emergency situation. Which of the following is the most likely cause?

A. Sympathetic over activity

B. Vagal stimulation

C. Venous air embolism

D. Carotid body stimulation

Cerebellum
Get the balance right



41. In which of the given cases is this procedure usually performed?

SLN Bx

A. MRM in node negative patients ×
B. MRM in node positive patients ×
C. BCT in node negative patients ✓
D. BCT in node positive patients

ALND



tc99-albumin
↓
Gamma camera
Methylene blue

×× ALND

A LND
(+) ← frozen sl^n ← LN

42. A 66-year-old woman with warfarin induced bleeding receives IV vitamin K, 4 U PRBCs and 4U FFP. 4h later, she begins to complain of shortness of breath. Vitals T - 39°C, RR 35/min, SpO2 84% on RA, and BP 88/52 mm Hg. Exam revealed B/L crackles and the patient was intubated for respiratory failure. CXR is as shown in the picture. What is the likely diagnosis?

A. Anaphylaxis

B. TACO

C. TRALI

D. Aspiration



43. Which of the following is not a correct match with respect to the mechanism of action and clinical use of the given drugs?

A. Zavegepant - CGRP receptor antagonist used for the treatment of acute migraine attacks

B. Sparsentan - Dual endothelin and angiotensin receptor antagonist used for the treatment of CHF

IgA nephritis

C. Lenacapavir - HIV-1 capsid inhibitor used for the treatment of HIV infection

D. Teplizumab - Anti-CD3 monoclonal antibody used for the prevention of type 1 diabetes in high-risk individuals

Cerebellum
Get the balance right

NEW DRUGS

- Daprodustat: HIF-PH inhibitor: Anemia due to chronic kidney disease
- Elacestrant: SERD (degrader): Hormone receptor-positive breast cancer
- Tirzepatide: Dual GLP-1 and GIP receptor agonist-Type 2 diabetes
- Mavacamten: Cardiac myosin inhibitor-HOCM

Cerebellum
Get the balance right

44. You are preparing to intubate a patient in the operating room. As part of your preoperative assessment, you perform a Mallampati examination to assess the patient's airway. Identify the class from the image shown below?

A. Mallampati Class I

B. Mallampati Class II

C. Mallampati Class III

D. Mallampati Class IV

45. A 4-year-old boy is brought to the emergency department for difficulty breathing and a productive cough. He has a past medical history of recurrent episodes of sinusitis and otitis media. The respiratory infections typically have a protracted course but respond to high-dose antibiotic therapy. Pulmonary auscultation shows crackles and wheezing. Cardiovascular examination shows a cardiac point of maximal impulse that is palpated at the right 5th intercostal space. Which of the following is most likely to be dysfunctional in this patient?

dextrocardia

A. C1 complement component

B. Chloride channels ← CF

C. Dynein arms ← PCD – Kartagener S.

D. IL-2 receptors

46. Correct about Brown-Sequard syndrome:

A. Extensive loss of pain on same side below the level of hemisection

B. Loss of fine touch and vibration on opposite side below the level of lesion

C. Extensive paralysis of muscle on opposite side below the level of lesion

D. Complete LMN type paralysis on same side at the level of lesion



47. A 2-year-child presented with complaints of abdominal pain following which he develops altered sensorium. He lives in an old house with paint chipping off. The following appearance is noted on X-rays. Which enzyme deficiency is likely in this child causing all these symptoms?

A. ALA synthase ← X linked
B. Heme synthase X
C. ALA dehydratase
D. Coproporphyrinogen synthase X



Cerebellum
Get the balance right

48. All are true about the muscle whose function is tested in the below shown test except:

A. Inserts into the lesser tubercle of the humerus

B. Primary function is ~~external rotation~~

C. Arises from the subscapularis fossa

D. Supplied by a branch of the posterior cord of the brachial plexus

addn / IR

U/L

Subscap N

Lift off → Subscapularis

49. A 45-year-old male with a history of human immunodeficiency virus (HIV) presents to the clinic with complaints of fatigue, weakness, and shortness of breath. The patient reports taking his anti-HIV medications consistently, which includes zidovudine (AZT). A complete blood count reveals a hemoglobin level of 8.5 g/dL, and a peripheral blood smear shows macrocytic anemia. Which of the following adverse effects of zidovudine is most likely causing the patient's symptoms?

Cerebellum
Get the balance right

A. Lactic acidosis
B. Peripheral neuropathy
C. Pancreatitis
D. Bone marrow suppression

50. You are the team leader during a cardiac arrest resuscitation scenario in a hospital setting. A 65-year-old patient has suddenly collapsed and is unresponsive. The monitor displays a rhythm as shown in the image. What is the next immediate step?

A. Administer epinephrine

B. Begin high-quality CPR

C. Defibrillate at 360 joules

D. Administer MgSO4



51. A 24 year old G2 at 28 weeks is brought to the emergency after being found unconscious at home by her mother. She is disoriented on arrival and her BP is 180/110 mmHg and 4 + proteinuria. On arrival at the emergency, she has a tonic-clonic seizure which resolves with MgSO4 administration. Once the patient is stabilized, she undergoes cesarean delivery for fetal distress. Post-partum, the below MRI is done. What is the most likely finding on imaging?



A. Subdural hematoma
B. Parenchymal hemorrhage
C. Subarachnoid hemorrhage
D. Posterior reversible encephalopathy syndrome

PRES

Hyper

post circul

52. A screening test is positive in 50% diseased population and 10% healthy population. Calculate the PPV:

A. 83
B. 90
C. 50
D. 89



$$PPV = \frac{50}{60} \times 100 = 83$$

Cerebellum
Get the balance right

53. A 63-year woman old presents to the AIIMS emergency chest pain after an RTA. Which resulted in crushed left lower limb. ECG shows 1-2 mm ST elevations in leads V2-V4 & troponin are elevated. The patient undergoes CAG which was normal. Which of the following is likely?

A. Aortic Dissection

B. Ramus intermedius

C. Severe mitral regurgitation with papillary muscle dysfunction

D. Akinetic LV apex with ballooning

STEMI - angiography

Takutsubo stress

Cerebellum

54. 6-year-old girl is brought to the office for evaluation of short stature. Her parents have noticed that she bears little resemblance to her 2 older siblings. The patient's mother and father are 172.5 cm (5'8") and 183 cm (6') tall, respectively. On physical examination, the patient's height is 3 standard deviations below the mean for her age. Other findings include low-set ears, a high arched palate a webbed neck, and cubitus valgus. Chromosomal analysis reveals a 45,XO karyotype. The patient is started on a medication to improve growth and normalize her height. Which of the following intracellular pathways is stimulated by the medication used in this patient?

GH → PIGGLET

A. Binding of activated receptors to DNA to modify transcription

B. Cyclic AMP-Protein kinase A pathway

C. Inositol triphosphate calcium-calmodulin pathway

D. JAK-STAT pathway

Cerebellum
Get the balance right

55. A 40-year-old woman has noted painless enlargement of her neck for the past month. On physical examination, she has diffuse, symmetrical thyroid enlargement without tenderness on palpation. Laboratory studies show her serum TSH is elevated but serum thyroxine is lower side of normal limits. Her microscopic examination is shown here. Which of the following laboratory test findings is she most likely to have?

Hypothyroid

Hashimoto

A. Increased serum calcitonin

B. High anti-thyroid peroxidase antibody titre

C. Increased serum parathormone

D. Positive antinuclear antibody test



56. A 23-year-old medical student presents to the surgery emergency department after cutting her wrist with a blade. After receiving first aid, the resident notices multiple healed cut marks on her wrist. Upon psychiatry referral, the patient informs the resident that she cut herself because she was feeling very angry at her boyfriend who spent the entire evening watching an IPL match rather than calling her. She also mentions that she has never been in a relationship with a guy for more than 2 months and believes that all men are horrible, while women are the best. What is the most likely diagnosis for this patient?

spelling

A. Histrionic Personality Disorder ××

B. Borderline Personality Disorder → Dialectical BT

C. Narcissistic Personality Disorder ××

D. Dependent Personality Disorder

57. All are true about the part labelled here except:

A. Derived from a type of atavistic epiphyses
B. Gives insertion to pectoralis major muscle
C. Short head of biceps takes origin from here
D. Can be felt in the infraclavicular fossa on examination



Cerebellum
Get the balance right

58. Identify the correct statements:

1. Grade 2 disability per 10 lac population is the most important leprosy awareness indicator

2. VVM shows cumulative exposure of the vaccine to the heat and assess vaccine potency

3. As per the National Tuberculosis Elimination Program (NTEP), HIV-TB prevalence of district >15% is a high-priority district

4. The external validity of a screening test is related to generalizability

Options:

A. 1,2,3,4

B. 1,2,4

C. 1,2,3

D. 2,3,4

59. A7-year-old girl is brought to the office due to a diffuse rash. The father says she developed a lace-like, erythematous rash this morning on her stomach, back, and extremities. Two days ago, the girl developed red, flushed cheeks. She also had congestion, headache, and a low-grade fever for 3 days earlier this week. The infectious agent most likely responsible for this patient's disease replicates in progenitors of which of the following cell lines?

Erythroblast

A. Erythrocytes

B. Hepatocytes

C. Intestinal epithelial cells

D. Lymphocytes

Cerebellum
Get the balance right

60. Anaerobic glycolysis of which of these produces 3 ATPs perunit glucose consumed?

A. Glucose
B. Fructose
C. Galactose
D. Glycogen



Cerebellum
Get the balance right

61. 44-year-old man comes to the OPD for follow-up of type 2 diabetes mellitus. He was diagnosed with diabetes 4 years ago and has been taking metformin since then. Six months ago, a new antidiabetic agent was added due to suboptimal glycemic control. Today, the patient is happy because he has lost 7 kg without any changes in diet or activity since the new therapy was started. His only symptoms are occasional nausea vomiting, and a prolonged sensation of fullness after eating. Laboratory testing reveals normal renal function studies and hemoglobin A1c of 7.2%. If this patient's weight loss is due to the new medication, which of the following agents is most likely responsible?

A. Basal-bolus insulin therapy
B. Thiazolidinediones
C. Glucagon-like peptide-1 agonist
D. Sulfonylurea

Cerebellum
Get the balance right

62. Identify the correct pair

1. Benefit measured in terms of natural units-Cost benefit analysis
2. Method based on behavioral sciences-MBO
3. PERT-Network Analysis
4. Benefit measured in terms of QALY-Cost effective analysis

Options;

A. 1,2,3

B. 2,3,4

C. 2,3

D. 1,4

Cerebellum
Get the balance right

63. A 20-year-old woman is brought to the emergency department by police at 2:30 AM after she was caught attempting to scale a fence at the Rashtrapati Bhawan. The patient appears highly agitated and paces around the examination room. She has just flown in from out of state to "meet with the president about a foolproof plan for eliminating worldwide terrorism." The patient has barely slept for the past week due to her intensive work on this plan. The evaluation has to be stopped when the patient begins banging on the door and demanding to leave. Temperature is 37 C (98.6 F), blood pressure is 148/84 mm Hg, pulse is 98/min, and respirations are 22/min. Urine drug screen is negative. Administration of which of the following medications is the most appropriate next step in management of this patient?

A. Lithium ✓
B. Clozapine ××
C. Haloperidol — psychosis
D. Olanzapine

Li + atypical AP = Mania

Cerebellum
Get the balance right

64. A newborn is being evaluated in the nursery. The patient was delivered to a 23-year-old woman. The mother developed dark terminal facial hair and a deepened voice during the pregnancy. The delivery was unremarkable, and the patient's vital signs are within normal limits. Examination shows ambiguous genitalia land clitoromegaly. Laboratory studies reveal elevated serum levels of testosterone and androstenedione. Karyotype testing shows a 46,XX genotype. Ultrasound of the pelvis shows a normal-sized uterus. This infant's presentation is most likely due to deficiency of which of the following enzymes?

A. 5-Alpha reductase

B. 17-Alpha hydroxylase

C. 21-Hydroxylase

D. Aromatase



65. Which of the following is included in high-quality CPR?

1. Compression depth of at least 5 cm
2. Rotate chest compressor every ~~3 mins~~ 2 min
3. Avoid excessive ventilation
4. Compression - ventilation ratio of 30:2
5. Allow complete recoil
6. ET co2 <10mm >10mm
7. Intra-arterial DBP >20mm Hg
8. Breaths 20-30/min x 10b/min

child

Cerebellum
Get the balance right

Options;

A. 1,2,3,4,5,6,7,8

B. 1,2,4,5,7,8

C. 1,3,4,5,7,8

D. 1,3,4,5,7

66. 38-year-old woman, gravida para 0, at 34 weeks gestation comes to the OPD due to increasing swelling in her hands and legs. The patient first noticed the swelling a few weeks ago but now cannot stand for more than an hour without needing to elevate her legs. She has no orthopnea, dyspnea, or chest pain. The patient has no chronic medical conditions and takes no daily medications. Vital signs are normal. Cardiac examination is within normal limits. Lungs are clear to auscultation. Bilateral lower extremities have 2+ pitting edema to the midshin. Urine dipstick shows no protein. Compared to a healthy nonpregnant woman, which of the following parameters is most likely to be decreased in this patient?

Cerebellum
Get the balance right

A. Cardiac preload

B. Cardiac output

C. Central venous pressure

D. Systemic vascular resistance

67. A 19-year-old man comes to the physician due to diarrhea for the last 4 months. He visited a student health clinic at his university a month ago and was prescribed a course of antibiotics, but his diarrhea failed to improve. He also has persistent nausea, a bloating sensation, and a 6.8-kg weight loss. Microscopic examination of his stool is shown in the image. Impairment of which of the following immune functions would most likely predispose to this patient's infection?

A. Bacterial killing by neutrophils

B. CD8+ T lymphocyte-mediated cytotoxicity

C. Eosinophil-mediated cytotoxicity

D. Secretory IgA production

Cerebellum
Get the balance right

68. All of the following structures lie in the relation to the left ureter except:

A. Mesentery of sigmoid colon
B. Bifurcation of common iliac artery
C. Quadratus lumborum
D. Gonadal vessels

Cerebellum
Get the balance right

Relations of Ureter

Anterior Relations

Right ureter

- Duodenum (2nd part)
- Right colic vessels
- Ileocolic vessels
- Right gonadal vessels
- Root of mesentery.

Left ureter

- Left colic vessels
- Sigmoidal vessels
- Left testicular or ovarian vessels
- Sigmoid mesocolon.

Posterior Relations

Both ureters run anterior to psoas major muscle and bifurcation of common iliac artery.

Cerebellum
Get the balance right

69. 65-year-old man presents with progressive dyspnoea & pulmonary function testing was done. FEV1 is 2.5 L (78% predicted), FVC is 4.00 L (94% predicted), and the FEV1/FVC ratio is 62.5%. The flow-volume curve is shown. What is the most likely cause?

A. Aspirated foreign body
B. U/L vocal cord paralysis
C. Subglottic stenosis
D. COPD



Cerebellum
Get the balance right

70. A 4-month-old girl is diagnosed with a febrile urinary tract infection. A voiding cystourethrogram is shown in the image. Which of the following is the most likely cause of this infant's presentation?

A. Congenital short intravesical ureter length

B. Damage to the rectovaginal septum

C. Obstruction due to a renal stone

D. Presence of a posterior urethral valve



Cerebellum
Get the balance right

71. Identify the correct kit for the clinical presentations:

1. Painful ulcers upon palpation, with suppurative inguinal lymphadenopathy on the right side → Chancroid – White
2. Vaginal discharge in a CSW with cervical erosions → Grey
3. Painful, swollen, and tender inguinal lymph node in a truck driver

Bubo - black

4. Painless ulcer in the genital area with non-tender inguinal lymphadenopathy in CSW; past h/o treatment of a similar lesion resulted in fever, malaise, tachycardia and vomiting – white

Syphilis

Jarisch - Herx

Cerebellum
Get the balance right

Options;

A. 1-Red, 2-Green, 3-Black, 4-Blue

B. 1-White, 2-Green, 3-Black, 4-Blue

C. 1-White, 2-Grey, 3-Black, 4-White

D. 1-Blue, 2-Grey, 3-White, 4-Blue

72. A 24y P2L2 who does not want more children presents to the OPD as she cannot feel the thread of the Multiload 375 that was inserted 3 years earlier. Her last menstrual period was 3 weeks earlier. On examination, the cervix appears normal, and the thread is not seen. A transvaginal scan shows the following image. What is the next best step?

→ Next: USG → No CoT → XRay

A. Hysteroscopic IUCD removal

B. Reassure her

C. Remove the IUCD with a hook

D. X-Ray Abdomen and Pelvis



CuT expelled

Thread → coiled/removed

CuT perforated outside peritoneum

73. A 45-year-old male presents with fatigue, night sweats, and weight loss. A complete blood count (CBC) is performed, and the peripheral blood smear shows the presence of cells with increased basophilic cytoplasm, granulocytic precursors, and a high percentage of blasts. A bone marrow biopsy is performed. Which of the following is the most appropriate investigation for this patient's condition?

A. Flow cytometry
B. Cytogenetics
C. FISH
D. Karyotyping



Cerebellum
Get the balance right

74. 37-year-old woman, gravida 2 para 1, at 30 weeks gestation comes to the OPD for a routine prenatal examination. Her pregnancy has been uncomplicated to date, and fetal movement is present. At the last visit, she drank a solution containing 50 g of glucose, and serum glucose was abnormally elevated an hour later. Which of the following hormones is most likely contributing to this patient's abnormal laboratory findings?

A. HCG
B. Human placental lactogen
C. Oxytocin
D. GH



Cerebellum
Get the balance right

75. 31-year-old woman comes to the office due to transient visual changes that are characterized by "loss of vision for a minute" and commonly precipitated by bending forward or lifting objects. During the last 3 months, she has had persistent headaches that she attributes to migraines. The patient takes over-the-counter analgesics as needed. Blood pressure is 140/90 mm Hg and pulse is 72/min and regular. BMI is 32.4 kg/m2. Funduscopic findings of the left eye are shown in the image below, with similar findings in the right eye. Which of the following is the most likely cause of the observed funduscopic findings in this patient?

A. Choroidal inflammation
B. Elevated intracranial pressure
C. Elevated intraocular pressure
D. Optic nerve inflammation

Cerebellum
Get the balance right

76. A 25-year-old woman comes to the emergency department due to 2 days of abdominal pain and bloody diarrhea. Temperature is 37.1 C (98.8 F), blood pressure is 119/76 mm Hg, pulse is 92/min, and respirations are 16/min. There is abdominal tenderness with no rebound or guarding. Guaiac-positive bloody stools are detected on rectal examination. Stool cultures reveal an Escherichia coli strain that does not produce glucuronidase and does not ferment sorbitol on sorbitol-containing MacConkey agar. Which of the following best describes the mechanism of action of the toxin specific to these bacteria?



A. Disruption of the cellular cytoskeleton

B. Inactivation of 60S ribosomal subunits

C. Inactivation of elongation factor 2

D. Inhibition of sodium reabsorption

77. Identify the true statements:

1. TxA2 has platelet aggregating and vasoconstricting effect
2. Lipoxin B4 inhibits inhibit neutrophil chemotaxis and adhesion to endothelium.
3. Il-1, Il-18, TNF are pyrogenic cytokines
4. C3 can be activated is both common as well as alternative pathways
5. T cells with a TCR composed of a γ and δ subunit are found associated with GI mucosa

Cerebellum
Get the balance right

Options;

A. 1, 2, 3, 4

B. 1, 2, 4, 5

C. 2, 3, 4, 5

D. 2, 4

78. 65-year-old woman is concerned about her risk of fracture as her mother was hospitalized for osteoporotic hip fracture in the past. She has a history of hypertension, hyperlipidemia, coronary artery disease, seizure disorder, and gastroesophageal reflux disease. The patient underwent menopause at age 52. She has smoked a pack of cigarettes daily for 24 years but does not drink alcohol. Weight is 56 kg. Long-term use of which of the following medications may increase this patient's risk of osteoporosis and hip fracture?

A. Atorvastatin

B. Levetiracetam

C. Metoprolol

D. Omeprazole r/o ##

Nov '14

Cerebellum
Get the balance right

79. Identify the nerve block given at the point shown:

A. Pudendal nerve — Ischial spine
B. Sciatic nerve
C. Obturator nerve
D. Inferior rectal nerve



80. 7-year-old boy is brought to the clinic for a routine health maintenance examination. He underwent a splenectomy a year ago for persistent anemia and jaundice. He has received the pneumococcal and meningococcal vaccinations and takes penicillin prophylaxis. Physical examination reveals a well-appearing child. The sclerae are anicteric. The abdomen is soft and nontender; no masses are palpable. Several well-healed laparoscopy scars are noted. Complete blood count is as follows:
Hemoglobin: 11.3 g/dL
Reticulocytes: 3%
Platelets: 160,000/mm³
Leukocytes: 8,200/mm³
Peripheral blood smear with Wright stain is shown.
This patient's findings are most likely related to which of the following?

HJ

A. Chronic Hb oxidation and precipitation
B. Inability to produce beta globin chains
C. Mechanical red-blood-cell membrane damage
D. Persistence of red blood cell precursor nuclei — nuclear remnants



81. A 30 year old man presented with profuse bleeding from the nose which did not stop on applying local pressure or vasoconstrictors. Anterior nasal packing was done but the bleeding could still not be controlled. It was decided to embolize one of the arteries supplying the vascular plexus responsible for most of the anterior nasal bleeds. Which of the following arteries should never be embolized?

Cerebellum
Get the balance right

A. Sphenopalatine artery
B. Septal branch of superior labial artery
C. Anterior ethmoidal artery ← ICA
D. Greater palatine artery

82. A 52-year-old woman comes to the emergency department due to recurrent right lower extremity swelling, pain, and erythema. The patient's medical history includes type 2 diabetes mellitus and depression. Her medications include metformin, paroxetine, and a multivitamin. The patient is diagnosed with cellulitis, admitted to the hospital, and started on antibiotics. She is continued on her home medications. Three days later, the patient becomes agitated and delirious with severe abdominal cramps and diarrhea. Her temperature is 39.2 C (102.6 F). blood pressure is 180/100 mm Hg, and heart rate is 120/min and regular. On examination, the patient is diaphoretic and tremulous, and her pupils are dilated. Bilateral hyperreflexia and ankle clonus are present. She begins to have seizures. Which of the following antibiotics was the patient most likely given to treat the cellulitis?

A. Clindamycin
B. Doxycycline
C. Linezolid
D. Vancomycin

83. Which of these is an example of anaplerotic reaction?

A. Pyruvate to Acetyl CoA
B. Pyruvate to oxaloacetate
C. Pyruvate to lactic acid
D. Pyruvate to acetaldehyde

Cerebellum
Get the balance right

84. Match the following drugs with their specific side effects:

A. Emtricitabine

B. Abacavir

C. Rosiglitazone

D. Liraglutide

E. Canagliflozin

F. Dabigatran

G. Warfarin

Options;

A. 1-B, 2-C, 3-F, 4-G

B. 1-A, 2-C, 3-E, 4-G

C. 1-G, 2-F, 3-E, 4-A

D. 1-E, 2-B, 3-C, 4-D

Cerebellum
Get the balance right

85. Identify the incorrect statements:

1. Superficial anterior compartment of leg has no neurovascular bundle
2. Tectorial membrane is the superior continuation of PLL
3. After parotidectomy, a person complains of loss of sensation in the shaving area. This is likely due to injury to auriculotemporal nerve.
4. Common carotid artery divides at the level of lower border of thyroid cartilage
5. Ansa cervicalis lies on the posterior wall of carotid sheath

A. 1,2,3,4,5
B. 1,4
C. 2,4,5
D. 1,3,4,5



86. A 70-year-old man, a retired banking executive with a history of smoking is admitted in ward with shortness of breath. Interpret the acid base disorder based on the arterial blood gas analysis report mentioned below. pH – 7.30, PCO2 -65 mm Hg, PO2 -60 mmHg, HCO3 - 36 mEq/L.

A. Metabolic acidosis
B. Respiratory acidosis
C. Metabolic alkalosis
D. Respiratory alkalosis

Cerebellum
Get the balance right

87. A medical student have given a topic to present in class. The topic given to you was excitation-contraction coupling. It is a process in which muscle cells are stimulated to contract and it is followed by muscle cell relaxation. It involves the following events:

1. Action potential
2. Calcium release into the cytoplasm
3. Cross-bridge cycling
4. Cytoplasmic calcium removal
5. Depolarization of T tubules

Which of the following options lists the correct sequence of events for excitation-contraction coupling in skeletal muscle tissue?

Cerebellum
Get the balance right

Options:

A. 1, 5, 2, 3, 4

B. 1, 2, 3, 4, 5

C. 1, 3, 4, 2, 5

D. 1, 4, 2, 5, 3

88. A female presents to the emergency 2 weeks after a normal vaginal delivery with abnormal involuntary flexure of the neck towards one side. History taking revealed that she was unable to lactate spontaneously for which she took a medication that she found on the internet without taking proper medical advice. Intake of which of the following drugs is most likely to lead to this condition?

A. Bromocriptine
B. Levodopa
C. Chlorpromazine
D. Benzhexol

89. All of the following are true:

1. Leptin acts via JAK-STAT signalling pathway
2. Damage to the Zona Occludens can result in increased permeability in Vibrio cholera diarrhea
3. Raised 2,3 BPG, Acidosis, HbF cause a right shift of the oxygen-hemoglobin dissociation curve
4. Macula densa is located at the Junction of thick ascending limb and DCT
5. The organ with the least arteriovenous pressure difference is the kidney.

Options;

A. 1, 2, 3, 4, 5

B. 2, 3, 4

C. 1, 2, 4, 5

D. 5

Cerebellum
Get the balance right

90. A 30-year-old female came to the OPD with complaints of backache. On examination, her eyes and pinna revealed blackish discoloration. What is the likely diagnosis?

Ochronosis

A. DISH
B. Fluorosis
C. Alkaptonuria
D. Ankylosing spondylitis



Cerebellum
Get the balance right

91. A 15-year-old male presents to emergency department in a comatose state. He was found in this state by his friend in the schoolyard. His friend confesses that the patient has been depressed over the past several days. His initial labs reveal hypocalcaemia and a severe metabolic acidosis. The urine shows calcium oxalate crystals. What was the toxin ingested?

A. Methanol

B. Ethylene Glycol

C. Organophosphates

D. Ethanol

Cerebellum
Get the balance right

92. A 64-year-old man is evaluated for persistent fever and weakness. He has a history of aortic valve replacement for aortic stenosis. Physical examination reveals a new cardiac murmur with scattered petechia and splinter hemorrhages seen on his extremities. Echocardiogram shows a vegetation involving one of the aortic valve leaflets, and blood cultures grow enterococci. As part of the patient's treatment, 240 mg of intravenous gentamicin is started. The pharmacy calculates that, in this patient, gentamicin has a volume of distribution of 30L, a half-life of 4 hours, and demonstrates first-order kinetics. Which of the following is the most likely serum drug concentration just before the next dose 8 hours later?

Vd: 30

t1/2 4hr

A. 0.5 mg/L

B. 1 mg/L

C. 1.5 mg/L

D. 2 mg/L

240/30 = 8 mg/L → (1/2) → 4 mg/L → (1/2) → 2 mg/L

93. Identify the correct statements

1. KEYNOTE-189 trial investigated NSCLC treatment using this drug that acts via pathway B.
2. Drug acting via pathway B has a role in treatment of ca endometrium.
3. Osimertinib is specifically used for treating NSCLC patients with T790M mutations and acts via pathway A.
4. Ipilimumab is a medication used in the management and treatment of metastatic melanoma that acta via pathway C.

A. 1, 2, 3, 4

B. 2, 4

C. 1, 2

D. 3, 4



94. What is the mechanism of action of the toxin produced by the bacterium shown in the figure below?

A. Increase in cAMP
B. Increase in cGMP
C. ADP-ribosylation of EF2
D. ADP-ribosylation of G protein



Cerebellum
Get the balance right

95. A neonate is lethargic, drowsy, and irritable with blood analysis showing hyperammonemia and orotic aciduria. Considering this patient, orotic acid levels are high due to which of the following?

A. Elevated ammonia
B. Elevated glutamine
C. Bypassing carbamoyl phosphate synthetase II (CPS-II)
D. Bypassing aspartate transcarbamoylase

✓ OTC


Mitochondria
CO_2 + NH_3
Carbamoyl phosphate synthetase 1 (CPS-1)
2ATP
2ADP
Carbamoyl phosphate
Ornithine transcarbamoylase (OTC)
Ornithine
Citrulline
OTC Deficiency and its effects
Glutamine + 2ATP + CO_2
CPS-II
RL
Carbamoyl phosphate
Carbomoyl Aspartate
Few steps
Orotic acid
Oritidine monophosphate (OMP)
Uridine Monophosphate (UMP)
Pyrimidines

96. Which of the following statements is/are true regarding anti-tubercular drugs?

1. Isoniazid inhibits mycolic acid synthesis in the cell wall of Mycobacterium tuberculosis.

2. Rifampicin is primarily used for the treatment of active tuberculosis and can cause liver toxicity and flu-like symptoms.

3. Bedaquiline is a second-line anti-tubercular drug that can cause QT prolongation and requires cardiac monitoring.

4. Pretomanid is primarily used as a monotherapy for the treatment of tuberculosis.

A. All options are correct
B. 1, 2 and 3 are correct
C. 1, 3 and 4 are correct
D. Only 1 and 2 are correct

Cerebellum
Get the balance right

97. A 42-year-old man is admitted to the hospital after sustaining fractures of his right femur and tibia in a motor vehicle collision. The patient becomes agitated, hypervigilant, and paranoid 12 hours after admission. He repeatedly tells the nurses that he can hear people in the corridor making insulting remarks about him. When hospital staff check the area near the patient's room, no one is found. Medical history is unremarkable, but psychiatric history is significant for cocaine, marijuana, and alcohol misuse. On examination, the patient is alert and oriented. Temperature is 37.2 C (99 F), blood pressure is 135/87 mm Hg pulse is 85/min, and respirations are 18/min. He is slightly tremulous and diaphoretic. Which of the following is the most likely diagnosis in this patient?

A. Alcoholic hallucinosis
B. Brief psychotic disorder
C. Cocaine withdrawal
D. Delirium tremens

Cerebellum
Get the balance right

98. The cranial nerve which is a content of the space labelled in the image below passes through which cranial foramina:

A. A

B. B

C. C

D. D





SOF
A.
B. R V2
C. O V3
D. S

FR → Pterygo palatine fossa → V2

JNA

99. 64-year-old man comes to the emergency department due to 2 days of fever, chills, productive cough. The patient has a history of hypertension, hyperlipidemia, and type 2 diabetes mellitus. Oxygen saturation is 94% on room air when lying on his right side but drops to 89% when he lies on his left. Dullness to percussion and bronchial breath sounds are present on the left side. Imaging is shown here. Which of the following pathophysiologic mechanisms in the left lung is most responsible for the drop in oxygen saturation after the change in position?



A. Dead space ventilation

B. Effort dependent hypoventilation

C. Hypoxic vasoconstriction

D. Intrapulmonary shunting

100. 46-year-old man is admitted to the hospital for elective cholecystectomy. He is started on intravenous fluids and antibiotics. Just before he is to go to an operating room, the patient complains of a sudden onset of chest tightness and light-headedness. His medical history is significant for anxiety, and he has had 2 previous visits to the emergency department for panic attacks. His blood pressure is 118/67 mm Hg. His electrocardiogram is shown here. Which of the following is the best initial step in treating this patient?



A. Adenosine
B. Alprazolam
C. Amiodarone
D. DC Cardioversion

unstable

101. Glycogen synthesis and breakdown takes place in the same cell, having enzymes necessary for both the pathways. Why the glucose-6-phosphate, freshly synthesized during glycogenesis in cytoplasm of hepatocytes, is not immediately degraded by the enzyme glucose-6-phosphatase?

A. The thermodynamics does not favor such a reaction to occur
B. Glucose-6-phosphatase is present in the endoplasmic reticulum and cannot act on glycogen formed in the cytoplasm
C. Glycogenesis and glycogenolysis are tightly regulated such that enzymes of only one of those are present at a time.
D. Steric hindrance due to albumin

102. A patient has the following laboratory findings:
Sodium – 138 mEq/L
Potassium – 4.4 mEq/L
Chloride – 112 mEq/L
Bicarbonate – 8.0 mEq/L
Arterial pH – 7.21



Which of the following is the most likely acid-base status?

A. Anion gap metabolic acidosis
B. Anion gap metabolic acidosis plus metabolic alkalosis
C. Anion gap metabolic acidosis plus normal anion gap metabolic acidosis
D. Normal anion gap metabolic acidosis

a) 10
b) 12
c) 18
d) 29

103. Healthy adult volunteers are enrolled in a phase I clinical trial investigating the properties of a newly developed oral antimicrobial agent. The drug is administered in different amounts to the volunteers over the course of several weeks to determine the best dosage that minimizes toxicity while maintaining trough levels above the minimum inhibitory concentration. While reviewing the data, the researchers note that the drug's half-life seems to vary amongst the study participants. An increase in which of the following pharmacologic parameters is most likely responsible for the longer half-life seen in certain individuals?

A. Drug glucuronidation
B. Glomerular filtration rate
C. Oral bioavailability
D. Volume of distribution



104. Identify the correct statements with respect to the deficits in different brain regions:

1. **Lateral geniculate body-Keyhole shaped defect**
2. **Substantia nigra-Paralysis agitans**
3. **Putamen-Athetosis**
4. **Hypothalamus-Hemiballismus**

A. 1,2,3

B. 2,3,4

C. 1,2

D. 3,4

Cerebellum
Get the balance right

105. A person presents with pneumonia. His sputum was sent for culture. The bacterium obtained showed gram positive cocci in chains on microscopy and alpha haemolytic colonies on sheep agar. Which of the following will help in confirming the diagnosis?

A. Novobiocin
B. Optochin
C. Bacitracin
D. Oxacillin

Cerebellum
Get the balance right

106. A 30y P1 with a last menstrual period 10 days ago presents to the emergency with complaints of worsening right pelvic pain. She says that she had sudden onset of sharp pain after bending over to pick up her child. The pain did not respond to acetaminophen and has intermittently worsened during the past several hours. She also had 2 episodes of vomiting. On examination, there is abdominal guarding. Transvaginal sonography demonstrates an 8-cm enlarged right ovary. There is high impedance arterial flow of the ovary on colour Doppler. These findings are most consistent with which of the following clinical diagnoses?

A. Appendicitis
B. Ovarian torsion
C. Haemorrhagic cyst
D. Ruptured corpus luteum cyst



107. Identify the true statements

1. In amyloidosis, Beta pleated sheet will be seen in X-ray crystallography.
2. There is decrease in thyroid-binding globulin, transferrin, and fibrinogen in nephrotic syndrome.
3. Blood transfusion requires screening of donor blood for HIV, HCV, HBV but not dengue.
4. Massive splenomegaly is noted in hairy cell leukaemia, CML and aplastic anemia.

A. 1, 2, 3, 4

B. 1, 3

C. 1, 2, 3

D. 2, 4

Cerebellum
Get the balance right

108. 46-year-old man comes to the OPD due to right shoulder pain. The pain started 2 months ago after the patient began working at a new warehouse where he frequently moves heavy boxes. On physical examination, external rotation of the right shoulder against resistance is painful, but there is no weakness. MRI reveals near complete tear of the right infraspinatus muscle. The absence of weakness in this patient is best explained by compensatory hypertrophy of which of the following muscles?

A. Latissimus-dorsi
B. Subscapularis
C. Supraspinatus
D. Teres minor

Cerebellum
Get the balance right

110. Match the following with the mode of cellular transport:

1. Oxygen	A. Simple diffusion
2. GLUT	B. Facilitated diffusion
3. SGLT	C. Primary active transport
4. Na+/Iodide symporter	D. Secondary active transport
5. Na+/K+ ATPase	
6. Proton pump	

Options:

A. 1-A, 2-B, 3-B, 4-C, 5- D, 6-D

B. 1-B, 2-A, 3-B, 4-C, 5- D, 6-D

C. 1-A, 2-B, 3-D, 4-D, 5- C, 6-C

D. 1-A, 2-B, 3-B, 4-D, 5- C, 6-C

Cerebellum
Get the balance right

111. Which of these represents true gastro-esophageal (GE) junction tumour?

A. Located between 5 cm and 1 cm above GE junction

B. Located between 2 cm above and 1 cm below GE junction

C. Located between 1 cm above and 2 cm below GE junction

D. Located between 1 cm above and 1 cm below GE junction

Cerebellum
Get the balance right


5 cm
Type I
Anatomical cardia
1 cm
Type II
–2 cm
Type III
–5 cm

112. Which of the following genetic defect regarding Maturity Onset Diabetes in the Young (MODY) is incorrectly matched?

A. MODY 2 – Glucokinase

B. MODY 3 – ~~HNF 4 alpha~~ HNF 1α – MC QQ

C. MODY 4 – IPF 1

D. MODY 5 – HNF-1 beta

Cerebellum
Get the balance right

113. 26-year-old woman is being evaluated for a possible inherited disorder. She has a 6-year history of generalized tonic-clonic seizures, and a year ago, she had partial loss of vision due to an occipital infarction. Her mother has chronic intermittent muscle weakness and lactic acidosis, and her maternal uncle has hemiplegia. After further evaluation, all the affected family members are found to suffer from the same inherited disease. Pedigree analysis is shown in the image below. Which of the following is the most likely explanation for the variability of clinical manifestations in the affected family members?

A. Anticipation
B. Heteroplasmy
C. Incomplete penetrance
D. Mosaicism



114. Which of the following statements is true about beta blockers:

1. **Carvedilol is the most widely used beta blocker.**
2. **These are drug of choice in acute decompensated heart failure.**
3. **Pindolol would be less likely to induce severe bradycardia in comparison to the other β-blockers.**
4. **Sotalol can prolong the AP duration leading to torsade's de pointes.**

A. 1, 2, 3, 4

B. 3, 4

C. 1, 3, 4

D. 2, 4

Cerebellum
Get the balance right

115. Which of the following muscle is supplied by the nucleus present beneath the area marked by the arrow in adjoining image?

A. Risorius
B. Lateral rectus
C. Inferior oblique
D. Medial pterygoid



Cerebellum
Get the balance right

116. 44-year-old man comes to the emergency department for exertional dyspnea, abdominal distention, and abdominal discomfort. He has a history of alcohol abuse, peptic ulcer disease, and acute pancreatitis. Physical examination shows a distended abdomen with shifting dullness and 1+ ankle edema. A diagnostic paracentesis is performed and shows the following:

Albumin-2.5 g/dL

Neutrophils-200/uL

Blood studies show the following:

Albumin-3.8 g/dL

Glucose-120 mg/dL

Aspartate aminotransferase-50 units/L

Alanine aminotransferase (ALT)-40 units/L

Which of the following is the most likely cause of this patient's ascites?

A. Decreased liver synthetic activity

B. Increased hepatic capillary hydrostatic pressure

C. GI malignancy

D. Pancreatitis

117. 56-year-old man is evaluated for dyspnea, wheezing, and cough productive of yellow sputum. The patient reports that he has been treated with antibiotics when the dyspnea is severe. He has smoked 2 packs of cigarettes daily for the past 25 years. The schematic diagram below is a representation of a bronchial wall. Which of the following ratios is the best pathological index to determine the severity of this patient's disease?

A. A/(A+B+C+D)

B. A/(A+B+C+D+E)

C. C/(B+C+D)

D. C/(A+B+C+D+E)



Cerebellum
Get the balance right

118. A group of investigators examined the effects of different muscarinic agonists and antagonists on the bladder musculature. The curves presented below were constructed after application of different agents. Curve A represents the effect of bethanechol, a pure cholinergic agonist. Which of the curves most likely represents the effect of bethanechol after application of a reversible competitive antagonist?

A. Curve B

B. Curve C

C. Curve D

D. Curve E



119. 63-year-old man is admitted to the hospital after sufering a cerebral infarction. A swallow study performed during the acute recovery period shows oropharyngeal dysphagia. Despite being maintained on the appropriate dietary restrictions, the patient experiences an episode of vomiting while lying on his back and subsequently develops pneumonia. Which of the following lung regions is most likely to be affected?

A. Anterior segment of the left upper lobe

B. Basilar segments of the right lower lobe

C. Lingular segments of the left lung

D. Posterior segment of the right upper lobe



120. Which of the following anticancer drugs act at the target shown in figure?

A. Erlotinib

B. Ribociclib

C. Dabrafenib

D. Nilotinib



121. A seventeen-year-old patient is brought to the emergency department having just ingested a large overdose of an antihistaminic drug. This drug is a weak base capable of entering most tissues including the brain. Which of the following statements is true:

A. Urinary excretion would be accelerated by administration of NaHCO3, an alkalinizing agent.

B. Urinary excretion would be accelerated by the administration of NH4Cl, an acidifying agent.

C. More of the drug would be ionized at blood pH than at stomach pH.

D. Absorption of the drug would be faster from the stomach than from the small intestine.

Cerebellum
Get the balance right

122. Identify the incorrect pair of contraceptive dosages

A. Ullipristal for emergency contraception-30mg single dose

B. LNG for emergency contraception 1.5mg two tablets repeated after 12hrs

C. DMPA 150mg im 3 monthly

D. Mirena 52mg LNG with 20ug/d release

Cerebellum
Get the balance right

123. This is a schematic diagram depicting body structure of which of these helminthes?

A. Loa loa
B. Wuchereria bancrofti
C. Brugia malayi
D. Onchocera volvulus



Cerebellum
Get the balance right

124. A 12-year-old child is brought to the emergency department after developing blurred vision and confusion. He was playing in the park and ate an unknown fruit. Temperature is 38.9 C (102 F), blood pressure is 110/76 mm Hg, pulse is 120/min, and respirations are 16/min. He is disoriented to time, place, and person. Physical examination shows flushed skin and dry oral mucosa. Both pupils are dilated and nonreactive to light and accommodation. Bowel sounds are decreased. Which of the following drugs can potentially reverse this patient's condition?

anticholinergic

A. Atropine
B. Naloxone
C. Physostigmine
D. Thiamine

Cerebellum
Get the balance right

125. 31-year-old woman is brought to the emergency department by her boyfriend due to confusion. The boyfriend says that the patient "seemed okay" when he left for work this morning, but when he returned in the evening, she was disoriented and "acting like a drunk person"; she had also vomited "all over the bathroom." She has a history of bipolar disorder. Medications include clonazepam, lithium, and quetiapine. Temperature is 37 C (98.6 F), blood pressure is 110/68 mm Hg, pulse is 92/min, and respirations are 16/min. On physical examination, the patient is somnolent and has frequent fasciculations. Pupils are equal and reactive. During evaluation, she has a brief episode of a generalized tonic-clonic seizure. Laboratory results are as follows:

Hemoglobin: 13 g/dL
Leukocytes: 9,000/mm³
Sodium: 136 mEq/L
Potassium: 3.8 mEq/L
Bicarbonate: 28 mEq/L
Creatinine: 1.2 mg/dL
Glucose: 120 mg/dL
Serum lithium 2.8 mEq/L

In addition to supportive measures, which of the following is the best next step in management of this patient?

A. Activated charcoal
B. Calcium gluconate
C. Gastric lavage
D. Hemodialysis

Cerebellum
Get the balance right

126. A 1-day old baby is evaluated in the emergency department due to poor feeding, repeated bilious vomiting and not passing meconium since birth. Abdominal examination reveals a distended abdomen. Imaging shows the following picture. Identify the condition which is unlikely in this patient?



A. Aganglionosis
B. Anorectal malformation
C. Meconium plug syndrome
D. Meconium ileus



127. Which of the following is likely with ascitic fluid protein 1.9 g/dL and a serum albumin ascites gradient (SAAG) of 1.4?

A. Alcoholic cirrhosis

B. Peritoneal malignancy

C. Constrictive pericarditis

D. Veno-occlusive disease

Cerebellum
Get the balance right

128. Which of the following drugs reversibly binds the site A?

A. Clopidogrel
B. Prasugrel
C. Ticagrelor
D. Tirofiban



129. In the histology section given, which marked area is involved in relieving pain in response to massage?

Gate theory

A. A ← SG

Substantia gelatinosa

B. B

C. C

D. D



130. A 65-year-old man comes to the emergency department due to severe lower abdominal pain and nausea. He also has not been able to urinate for the past 24 hours. The patient has a history of hypertension and benign prostatic hyperplasia. On examination, a large mass is palpable in the suprapubic area. Multiple attempts at urethral catheterization are unsuccessful, and an ultrasound-guided midline suprapubic cystostomy is planned. Besides the bladder wall, which of the following structures is most likely to be penetrated by the trocar and cannula during the procedure?

A. Anterior abdominal aponeurosis
B. Parietal peritoneum
C. Perineal membrane
D. Visceral peritoneum

Cerebellum
Get the balance right


Suprapubic abdominal wall
External oblique
Internal oblique
Transversus abdominis
Anterior abdominal aponeuroses
Skin
Camper fascia
Scarpa fascia
Investing fascia
Extraperitoneal fat
Transversalis fascia
Bladder wall

131. A 25-year-old male patient presented with sudden onset breathlessness and a sharp pain in his chest after a road traffic accident. His BP is 85/69mm Hg and HR is 110bpm. The chest X-ray is given below. What is the next step for this patient?

1. eFAST
2. Insert iv cannula and start iv fluids
3. Needle thoracostomy
4. ICD insertion
5. Thoracotomy
6. CECT chest





Options;

A. 1,2,3,4

B. 3,4

C. 2,3,4

D. 1,2,3,4,5,6

132. A female presents to OPD with 6 weeks of amenorrhea complaints of bleeding per vagina and mild abdominal pain. The urine pregnancy test is positive and hCG level is 2800 IU/L. On investigation, mass is seen on the left adnexa measuring 3 x 2 cm. She is hemodynamically stable. How will you manage this patient?

A. Oral methotrexate

B. Single-dose methotrexate injection im

C. Serial methotrexate + leucovorin rescue

D. Salpingectomy

133. A 14-year-old girl is brought to the emergency department following a suicide attempt. She swallowed several of her grandfather's theophylline tablets after an intense argument with her parents a few hours ago. The patient reports nausea and headache. Her temperature is 37 C (98.6 F), blood pressure is 114/76 mm Hg, pulse is 88/min, and respirations are 16/min. This patient should be carefully monitored for which of the following?

A. Hypoglycemia and complete heart block
B. Jaundice and elevated liver enzymes
C. Sedation and respiratory depression
D. Seizures and tachyarrhythmias



134. A 19-year-old sex worker was treated 2 days ago for presumed gonorrhea with ceftriaxone and azithromycin. She also has a palpable spleen. Laboratory examination is notable for a Hb of 5 g/dL with normal white blood cells and platelets. Peripheral blood smear shows an excess of spherocytes. Which of the following tests will most likely confirm the diagnosis?

A. ADAMTS-13 activity assay

B. Direct antiglobulin (Coombs) test

C. Flow cytometry

D. G6PD deficiency assay

Cerebellum
Get the balance right

135. A 47-year-old woman comes to the office due to fever, dysuria, and abdominal pain. The patient has had several urinary tract infections since her teens but has no other medical concerns. Physical examination shows mild suprapubic discomfort. Urinalysis demonstrates pyuria and many bacteria. CT scan of the abdomen is shown in the image below: Which of the following structures has most likely limited the proper ascent of the anomalous organ seen on CT scan?

A. Inferior mesenteric artery
B. Inferior vena cava
C. Persistent urachus
D. Superior mesenteric artery



Cerebellum
Get the balance right

136. Slit lamp ophthalmological examination of a patient revealed the following. All of the following can cause this except:

A. Quinidine
B. Amiodarone
C. Chloroquine
D. Indomethacin



Cerebellum
Get the balance right

137. Which among the following species is transmitted through the vector given in the image?

A. R.conorii
B. R.typhi
C. R.akari
D. R.prowazekii

louse
/
Epidem

Cerebellum
Get the balance right

138. A 75-year-old male after a fall in bathroom and the pelvic X-ray is shown below. What will be the position of the left lower limb?

A. Shortened and internally rotated

B. Shortened and abducted

C. Extended, abducted and externally rotated

D. Abducted and internally rotated

Cerebellum
Get the balance right

139. Area marked A in the given figure contain all the following, except:

A. Zonula occludens
B. Zonula adherens
C. Macula adherens
D. Gap junction



140. 58-year-old, previously healthy man is evaluated due to a 2-month history of early satiety, postprandial epigastric pain and vomiting and unintended weight loss. Physical examination shows cachexia and palpable epigastric mass. Upper gastrointestinal endoscopy reveals a large prepyloric mass causing gastric outlet obstruction; biopsy of the mass is consistent with poorly differentiated adenocarcinoma. Abdominal CT scan reveals thickening of the gastric wall with adjacent fat infiltration, enlarged lymph nodes, and multiple hepatic metastatic lesions. The disease is considered unresectable, and the patient undergoes laparoscopic gastrojejunostomy for palliation of gastric outlet obstruction. Postoperatively, he tolerates tube feeding but experiences worsening weakness. The patient also has frequent episodes of non-sustained ventricular tachycardia. Laboratory results are as follows:

	At admission	Postoperative day 2
Hemoglobin	9.9 g/dL	9.8 g /dL
Leukocytes	8,800/mm³	10,000/mm³
Serum chemistry		
Sodium	130 mEq/L	135 mEq/L
Potassium	3.7 mEq/L	2.8 mEq/L
Creatinine	1.1 mg/dL	0.8 mg/dL
Glucose	90 mg/dL	120mg/dl
Phosphorus	3.1 mg/dL	1.2mg/dl

KMP ↓

Which of the following is the most likely cause of this patient's worsening condition?

A. Hypermetabolic syndrome

B. Nutritional replenishment

Refeeding Sx

C. Paraneoplastic syndrome

D. Sepsis

141. A 40 year old male patient is being evaluated for a suspected brain tumor. A biopsy is obtained whose image has been shown below. Which of the following is the most probable diagnosis based upon the biopsy findings?

A. Pilocytic astrocytoma

B. Glioblastoma multiforme

C. Medulloblastoma

D. Oligodendroglioma

Fried egg



142. A 68-year-old man comes to the OPD for a follow-up appointment. He has a history of advanced melanoma that is unresectable and resistant to adjuvant regimens. The patient recently began receiving monoclonal antibody infusions for advanced melanoma. The monoclonal antibodies block a specific cell surface receptor found on T lymphocytes. As a result, T cells capable of recognizing tumor antigens have improved ability to destroy cancer cells. Which of the following cell surface receptors is most likely blocked by the treatment?

immune check pt ⊖

A. CCR5
B. CD4
C. CD19
D. PD-1


Ipilimumab
Atezolizumab
Durvalumab
Avelumab
Cemiplimab
Nivolumab
Pembrolizumab
T cell
MHC II
APC
CD28
B7
CTLA-4
Antigen
MHCI
TCR
Tumor cell
T cell activation
PD-1
PD-L1

143. 52-year-old man comes to the OPD for evaluation of hearing loss in the left ear. The patient works in sales and is frequently on the phone. Over the past several months, he has had difficulty hearing phone conversations in the left ear and must move the headset to the right. He has no pain or vertigo but sometimes feels off-balance when walking at night and holds onto walls. A tuning fork placed on the forehead is lateralised to the the right ear. Air conduction is greater than bone conduction bilaterally. There is decreased sensation to touch on the left side of the face. Which of the following is the most likely cause of this patient's symptoms?

Lt SNHL

Lt V CN / VII CN

A. Cholesteatoma

B. Herpes zoster oticus

C. C Ménière disease

D. Vestibular schwannoma

Cerebellum
Get the knowledge right

144. A 13 year old girl has complaints of recurrent nose bleeds and excessive loss of blood during menstruation. She appears pale. Laboratory studies reveals the following:.

Hemoglobin - 7 g/dL

Platelets - 2 lac/mm3

Bleeding time - Prolonged

PT - Normal

aPTT - Prolonged

Which of the following factors may be deficient in this case?

A. Factor IX

B. Factor XI

C. gp 1b-IX

D. vWF

Cerebellum

145. Identify the incorrect statements:

1. Richardson and Koeppe are direct gonioscopes.
2. Narrowing of isopters and the appearance of a blockage within the blind spot are early changes in ~~MS~~ Glaucoma
3. Most common ocular manifestation of COVID-19 is ~~papillary~~ follicular conjunctivitis
4. The Haigis-L regression formula is used for lens power calculation in patients undergoing cataract surgery with a past history of LASIK for myopia.
5. Correct method is to put the ~~eye ointment first, drops 15 min later~~ drops → ointment

Cerebellum
Get the balance right

A. 1,2,3,4,5
B. 2,3,4
C. 1,3,5
D. 2,3,5

146. Identify the scheme which bears the logo given below

A. Janani Suraksha yojna

B. Janani Shishu Suraksha Karyakram

C. Labour Room Quality Improvement Initiative

D. Vandemataram Scheme

Cerebellum
Get the balance right

147. Identify the correct pair:

A. Staghorn calculus
B. Putty kidney
C. PUJ obstruction
D. Cortical nephrocalcinosis

A.



B.

C.



D.

148. The following image depicts an organic poison. Which of the following statements is true about it?

A. Fatal dose is 60-120 seeds (1 seed)

B. Weight of one seed is approximately ~~10g~~ and was used to measure gold in the past. 1g / seed

C. Symptoms resemble those of ~~cobra bite~~ viper bite.

D. Active principle is a toxalbumin.

Abrus

Cerebellum
Get the balance right

149. All of the following are true except:

A. Dermis has dense irregular connective tissue
B. Fibrocartilage, bones, tendons have collagen type I
C. Hyaline cartilage has perichondrium with 2–4 chondrocytes in each lacuna
D. Fibres from Inferior olivary nucleus directly reach the purkinje cells in cerebellum, which constitute the largest neurons in human brain



150. You are a medical officer posted in the one stop center of your hospital. You are required to collect evidences from a sexual assault victim. What is the correct order of collection according to the instructions given in the SAFE kit?.1- Cervical swab for culture.2- Vaginal swab.3- Clothing.4- Breast swabs.5- Nail clippings.6- Matted pubic hair cutting.7- Pubic hair combing

A. 3>5>4>6>7>1>2

B. 3>5>4>7>6>2>1

C. 4>5>3>6>7>1>2

D. 3>4>5>76>7>1>2

Cerebellum
Get the balance right

SAFE kit (sexual assault forensic evidence)

- Step 1- informed consent
- Step 2- history of sexual violence
- Step 3A- outer clothing.● Step 3B- inner clothing
- Step 4A- Debris from hair combing.● Step 4B- Interdigital debris.● Step 4C- Swabs from stain on the body.● Step 4D- Nail scrapings.● Step 4E- Nail clippings.● Step 4F- Scalp hair
- Step 5- Two breast swabs.
- Step 6A- Pubic hair combing.● Step 6B- Cutting of public hair.● Step 6C- Cutting of matted pubic hair.
- Step 7A- Two high vaginal/ cervical swabs.● Step 7B- Two low vaginal swabs.● Step 7C- Two urethral swabs.
- Step 8- Cervical swab for culture.
- Step 9- Vaginal washing.
- Step 10A- Two anal swabs.● Step 10B- Two rectal swabs.
- Step 11- Two oral swabs.
- Step 12- Blood samples for blood grouping, testing for alcohol or intoxication, DNA analysis, testing for HIV, VDRL, HbsAg.
- Step 13- Urine test for pregnancy and drugs.
- Step 14- Medical examination report

Cerebellum
Get the balance right

151. Identify the correct statements:

1. Miami criteria: 50% reduction in PTH within 10 mins of gland removal
2. Majority of the retrosternal goitres derive their blood supply from mediastinal vessels
3. Purulent peritonitis following a perforated diverticulitis is Hinchey grade 3
4. CECT abdomen in stable patients with eFAST positive is a part of "C" of primary survey

A. 1,2,3

B. 1,2,4

C. 1,3

D. 1,3,4

Cerebellum
Get the balance right

152. A patient has undergone cataract surgery for lens removal and IOL has not been implanted. Instead he is prescribed high power spectacles. Apart from cosmetic blemishes, what other side effects should he not be worried about ?

A. Pin cushion effect
B. Minification of image - Magnification
C. Roving ring scotoma
D. Jack in the box phenomenon

Cerebellum
Get the balance right

153. A 7 year old male patient is taken for right side radius-ulna open reduction internal fixation surgery under General anesthesia. As the child seems uncooperative for regional anesthesia, you decide to start general anesthesia induction with an inhalational agent via mask. Within a few minutes the patient starts breath holding and coughing. To find out the cause you check the inhalational agent vaporizer and you immediately turn it off and switch on the non- irritant Inhalational agent. What was the probable Inhalational agent which led to coughing and breath holding?

A. Xenon

B. Desflurane

C. Halothane

D. Sevoflurane

154. A 46 year old male presented to the clinic with a persistent cough, fatigue, weight loss, and night sweats. Xray of the patient is as given below. What is the most likely diagnosis?

A. Streptococcus pneumoniae infection
B. Mycobacterium tuberculosis infection
C. Carcinoma lung
D. Sarcoidosis



Cerebellum
Get the balance right

155. A 52-year-old man has undergone lung transplantation. Two months after receiving lung transplant, he developed pulmonary symptoms and was diagnosed as having bilateral diffuse interstitial pulmonary pneumonitis and bronchiolitis. Which of the following etiological agent is responsible in this condition?

A. Cytomegalovirus
B. Herpes Simplex virus
C. JC virus
D. BK virus



156. A 3-year-old girl is brought to the office after developing fever and a sore throat. The patient recently entered day care, and similar symptoms have been reported in several of the other children. Physical examination shows exudative pharyngitis and enlarged anterior cervical lymph nodes. A rapid antigen detection test confirms the diagnosis of streptococcal throat infection. Her condition resolves with antibiotic therapy. Several weeks later, she is re-exposed to Streptococcus pyogenes. The bacteria penetrating beyond the surface epithelium are immediately coated with preformed IgG antibodies. Which of the following substances acts in the most similar manner to IgG antibodies to facilitate phagocytosis?

A. Complement C3b

B. Complement C5a

C. Immunoglobulin M

D. Leukotriene B4

157. Following road traffic accident patient suffers polytrauma and is evaluated in the emergency section of the hospital. His pulse rate is 116, respiratory rate is 24, blood pressure of 122/78 mm of Hg and patient is mildly anxious. What is the approximate blood loss patient has following trauma?

A. 750-1500 mL

B. <750mL

C. 1500-2000ml

D. >2000 mL

Cerebellum
Get the basics right

158. Identify the accessory protein marked in the diagram below

A. Nebulin
B. Alpha actinin
C. Titin
D. Tropomyosin



Cerebellum
Get the balance right

159. Calculate the corrected reticulocyte count of a patient with haemoglobin of 5 and absolute reticulocyte count of 9%?

A. 4.5%

B. 6%

C. 1%

D. 3%

$$CRC = \text{Retic count} \times \frac{\text{Hb pt}}{\text{N Hb}} \times \frac{\text{Hct pt}}{\text{Hct N}}$$

15

45

$$= \frac{9 \times 5}{15} = 3$$

Cerebellum
Get the balance right

160. A research laboratory is evaluating a new dipstick test for diagnosing urinary tract infections (UTIs). The new test is compared to urinalysis of a midstream urine specimen (considered the diagnostic gold standard) to establish diagnostic test parameters. The study enrolls 300 patients, of which 100 have a UTI as determined by urinalysis. The new dipstick test is determined to be 70% sensitive and 90% specific for the diagnosis of UTIs. How many false positives are present in the study?

A. 20

B. 30

C. 70

D. 120



Cerebellum
Get the balance right

161. A 52-year-old female patient present with 4 cm diameter diagnosed breast cancer lesion with ipsilateral axillary and contralateral supraclavicular lymphadenopathy as shown below. As per AJCC system, patient belongs to which stage of breast cancer?

A. T4a N3c M1
B. T4b N1 M1
C. T4b N3c M0
D. T4c N1 M0



Cerebellum
Get the balance right

162. A woman presented to the clinic with heavy, painful menses that have worsened over the past year. On pelvic examination the uterus is enlarged and tender. MRI scan of the patient's abdomen revealed a finding similar to the image given below. What is the most likely diagnosis?

A. Polyps
B. Leiomyoma
C. Adenomyosis
D. Endometrial hyperplasia

Cerebellum
Get the balance right

163. A 29 year old female presented with complaints of secondary amenorrhea for the past 6 months. She reports having a regular menstrual cycle in the past and there were no recent changes in weight, exercise, or stress. She does not use any contraceptives and has no significant medical history. On examination, she appears well nourished. Her pregnancy test is negative. What is the next best step in management?

A. Measurement of serum prolactin levels
B. Measurement of serum FSH and LH levels
C. Pelvic USG
D. Endometrial biopsy



164. A patient with metastatic prostate cancer develops hypocalcemia during treatment with a bone antiresorptive agent. Which of the following agents is most likely to result in hypocalcemia?

A. Pamidronate
B. Zoledronic acid
C. Denosumab
D. Ibandronate

Cerebellum
Get the balance right

165. Which of the following options is incorrectly matched?

1. Prolapse in pregnancy: Ring Pessary
2. Vault prolapse - Sacrospinous ligament suspension
3. Cystocele - Colposuspension
4. Rectocele - Posterior colporrhaphy
5. Prolapse in a female who desires future child bearing-Shirodkar sling surgery
6. Prolapse in a 65 year old female with DM and hypertension-Le Fort's colpocleisis

A. 4,5
B. 3,6
C. 2
D. 3

166. This device shown in the image is used in:

A. Men with suspected urethral stricture
B. Women with recurrent UTI
C. Urine incontinence in pediatric population
D. Suspected VUR

167. A large, multi-country study is conducted to determine the effect of economic development on cancer incidence and mortality. The study uses data obtained from the national cancer registries, along with information regarding per capita gross domestic product as reported by the International Monetary Fund and life expectancy as reported by the World Health Organization. Which of the following best describes the design of this study?

A. Cross-sectional survey

B. Ecological study

C. Nested case-control study

D. Systematic review → Studies

Cerebellum
Get the balance right

168. 23-year-old comes to the ED with the feeling of food being stuck in the throat. Laryngoscopy reveals fish bone lodged in the left piriform recess. During retrieval of the fish bone, a nerve deep to the mucosa overlying the recess is damaged. Which of the following is most likely to be impaired in this patient as a result of his iatrogenic injury?

A. Cough reflex
B. Gag reflex
C. Mastication
D. Salivation



Cerebellum
Get the balance right

169. Match the following images:

A. Venous ulcer
B. Marjolin ulcer
C. Arterial ulcer
D. Trophic ulcer

1.

2.

3.

4.

A. 1-C, 2-D, 3-B, 4-A
B. 1-D, 2-C, 3-A, 4-B
C. 1-C, 2-D, 3-A, 4-B
D. 1-D, 2-C, 3-B, 4-A

Cerebellum
Get the balance right

170. The following hematoxylin and eosin-stained biopsy is from which tissue?

A. Tonsils
B. Spleen
C. Lymph node
D. Peyer's patches



171. Identify the correct scenario

1. Loss of sensations over the root of penis after lap hernia repair-Ilioinguinal nerve

2. Hyperesthesia of tongue after parotid surgery-Auriculotemporal nerve

3. Diaphragm palsy after thymoma resection-Phrenic nerve

4. Loss of sensations over the lateral gluteal region after hernia mesh repair-LCNT

A. 1,2,3,4

B. 1,2,3

C. 1,3

D. 2,3,4

172. A therapist observes distinct patterns in her sessions with a 35-year-old woman who has a turbulent history with relationships. Initially, the woman sings praises about the therapist, declaring her as the best thing that's happened to her. However, after a few sessions, she arrives agitated, accusing the therapist of not genuinely wanting to help her. These extreme shifts in perception occur frequently, creating a turbulent therapeutic environment. Which defense mechanism is predominantly being exhibited by this patient?

Cerebellum
Get the balance right

A. Reaction formation
B. Denial
C. Projection
D. Splitting

173. The anesthetist feels a snap while performing an epidural catheter insertion. This is due to piercing of which structure?

A. Ligamentum flavum
B. Supraspinous ligament
C. Dura mater
D. Posterior longitudinal ligament

Cerebellum
Get the balance right

174. A 32-year-old female presents to the dermatology clinic concerned about white patches on her skin that she noticed since childhood. Physical examination reveals depigmented patches with a symmetrical distribution, especially prominent on the ventral aspect of her body and mid-forehead. White forelock is noted. She mentions that her grandfather had similar patches since his birth. She also reports that a cousin on her father's side has been recently diagnosed with rheumatoid arthritis. Given the patient's history and clinical presentation, which of the following describes the underlying cause of her condition?

Piebaldism

A. A polygenic disorder with progressive depigmentation.
B. An autosomal dominant disorder involving the C-kit protooncogene.
C. An autoimmune condition with progressive depigmentation.
D. A sporadic condition without a genetic link

Cerebellum
Get the balance right

175. A 73-year-old man comes to the emergency department with unstable angina. He undergoes coronary angiography via the femoral approach. A stent is placed in the right coronary artery, and the patient is discharged. He returns to the emergency department 2 days later with blue discoloration of his right toe. He has pain and mild tingling in the affected toe. On physical examination, the right toe appears cyanotic, and there is livedo reticularis affecting the right thigh. Serum creatinine is 2.8 mg/dL (preoperatively it was 1.0 mg/dL). Which of the following histopathologic findings would most likely be seen on biopsy of this patient's kidney?

A. Cholesterol clefts in arterial lumen

B. Glomerular crescent formation

C. Extensive necrosis of proximal tubular cells

D. Tubular obstruction with rate crystals

Cerebellum
Get the balance right

176. A 60 year old male with history of long standing goitre visits the OPD. Surgeon performs this manoeuver- which makes the patient breathless. This finding implies?

A. Goitre is retrosternal
B. Malignant transformation
C. Post operative airway risk
D. Infiltration of carotid artery

Kocher test

Cerebellum
Get the balance right

177. In which of the following study designs, the Outcome of interest has already taken place when the study is started?

A. Prospective cohort study
B. Retrospective cohort study
C. Nested case control study
D. Both (b) and (c)



178. A surveillance study is conducted to assess the long-term efficacy and safety of a drug currently being used to treat patients with heart failure. Researchers enroll 8,300 patients with heart failure. The patients receive the drug once daily for 6 months. The results show significant clinical improvement, but severe hypernatremia is observed in 23 patients. The study publication recommends a lower dose of the drug in patients with baseline normonatremia and hypokalemia to prevent hypernatremia. Which of the following best characterizes this type of study?

Cerebellum
Get the balance right

A. Phase I clinical trial
B. Phase II clinical trial
C. Phase III clinical trial
D. Phase IV clinical trial

179. A patient presents to the emergency room with symptoms of fever, increased respiratory rate, elevated heart rate, and low blood pressure. The physician suspects the patient may have SIRS. Which of the following is not a criteria for SIRS?

White RR

A. Systolic blood pressure <90 mmHg
B. Temperature >38 degree Celsius or <36 degree Celsius
C. Respiratory rate >20 bpm
D. Heart rate >90 min

180. A 23-year-old man with HL was treated successfully with the ABVD regimen. Which of the following drugs used in the treatment of this patient is cell cycle specific ?

A. Alkylating agents
B. Corticosteroids
C. Antimetabolites
D. Plant alkaloids




Bleomycin
Microtubule inhibitors
Taxanes
Vinca alkaloids
Topoisomerase inhibitors
Etoposide
Teniposide
Irinotecan
Topotecan
Antimetabolites
Cladribine*
Cytarabine
5-flurouracil
Hydroxyurea
Methotrexate
Pentostatin
Thiopurines
*Cell cycle Nonspecific
Cell cycle-independent drugs
Platinum compounds
Alkylating agents:
Anthracyclines
Busulfan
Dactinomycin
Nitrogen mustards
Nitrosoureas
Procarbazine
G1
Double check repair
Mitosis
M
Cytokinesis
Interphase
DNA synthesis
Duplicate cellular content
S
G1
G0
Resting
Rb, p53 modulate G restriction point

181. A patient is induced with Thiopentone IV through one of the antecubital veins. The patient complains of severe pain engulfing his whole hand. What is the next line of management?

A. Give IV propofol through same needle

B. Insert another iv line immediately

C. IV lignocaine through same needle

D. Remove the needle immediately



182. A 60-year-old male is brought to the emergency department by ambulance after a road traffic accident. He doesn't know where he is, he is unable to identify his relatives and is in a state of confusion. On examination, he has eye-opening only on painful stimuli and can localize the site of painful stimuli. What is the GCS score of this patient?

A. E2V3M5

B. E2V4M5

C. E3V3M6

D. E3V4M6



183. A 48-year-old male presented to the clinic with complaints of acute abdominal pain, nausea, and vomiting. The pain is sudden onset, and severe, in the epigastric and umbilical regions, referred to the back and relieved on bending forwards. He has a past history of chronic alcoholism. On examination, the abdomen is tender with guarding and rigidity, rebound tenderness is present and mild jaundice is evident. Which of the following is the best investigation for this patient?

A. Liver function test

B. Serum amylase

C. USG abdomen

D. CECT scan

184. A 32-year-old female presented to the clinic with complaints of heart racing, diarrhea and intolerance to hot weather. She has a known history of iodine deficiency goiter and is taking iodine supplementations. She has no history of flu-like illness in the past weeks. Her blood results on the recent follow-up are as follows: high T3, high T4 and low TSH. What is the most likely diagnosis?

A. De Quervain thyroiditis

B. Wolf-Chaikoff effect

C. Jod-Basedow effect

D. Hypothyroidism

185. Identify the marked layer:

A. Outer plexiform layer
B. Inner plexiform layer
C. External limiting membrane
D. Internal limiting membrane



Cerebellum
Get the balance right

186. Identify the incorrect pair

A.

B.

C.

D.

A. C4c
B. C1
C. C4a xx C5
D. C4b

Cerebellum
Get the balance right

187. A group of sports physicians plans to conduct a case-control study to investigate a possible association between adolescent idiopathic scoliosis (AIS) and sacroiliac joint (SIJ) dysfunction in young athletes. The case group will consist of young athletes who were diagnosed with AIS during a regular checkup by a sports physician. Which of the following is the most appropriate control group for this study?

A. Young athletes with a diagnosis of AIS and SIJ

B. Young athletes with a diagnosis of AIS but not of SIJ

C. Young athletes with a diagnosis of AIS irrespective of SIJ status

D. Young athletes with no diagnosis of AIS irrespective of SIJ status

188. Identify the correct statements:

1. Thelarche is the first sign of puberty in females.
2. Maximum growth occurs in Tanner stages 4 and 5 in males
3. Penile enlargement is the first sign of puberty in males
4. Peak growth velocity always precedes menarche in females

A. 1,2,4

B. 1,2,3,4

C. 2,3,4

D. 1,2,3

Cerebellum
Get the balance right

Match the following images with the correct name of the manouever

A. Pawlick grip
B. Pelvic grip
C. Rubin manoeuvre
D. McRobert manoeuvre
E. Ritgen manoeuvre
F. Burns Marshall manoeuvre
G. Mauriceau–Smellie–Veit maneuver



1

2

a) 1-D, 2-B, 3-G, 4-F
b) 1-D,2-A, 3-E, 4-F
c) 1-C,2-A, 3-G, 4-C
d) 1-G,2-B, 3-C, 4-G

3



4

Cerebellum
Get the balance right

190. Correctly identify the type of suturing shown below.

A. Subcuticular sutures
B. Interrupted mattress sutures
C. Interrupted simple sutures
D. Continuous sutures



Cerebellum
Get the balance right

191. Identify the correct statements:

1. Social smile – 2months
2. Walk without support -~~12months~~ 15 mon
3. Babbling gibberish-15months
4. Walk upstairs with alternate foot – ~~4yrs~~ 3yr
5. Copy a circle-~~2yrs~~ 3y



A. 1,2,3,4

B. 1,3,4,5

C. 1,3

D. 1,3,5

Cerebellum
Get the balance right

192. A 65-year-old male with a history of atrial fibrillation and recent coronary artery bypass graft surgery presents with bleeding from the surgical wound site. Laboratory tests reveal a normal platelet count and coagulation panel. Thromboelastography (TEG) is ordered to assess the patient's clotting profile. Which of the following TEG tracings is most consistent with the patient's presentation?

DIC

A. Reduced R-time, increased angle, increased MA, increased LY30

B. Prolonged R-time, decreased angle, decreased MA, decreased LY30

C. Prolonged R-time, increased angle, decreased MA, increased LY30

D. Reduced R-time, decreased angle, increased MA, decreased LY30

TEG	ROTEM	Coagulopathy	Treatment options
R 10–14 min	ExTEM CT 80–100 s InTEM CT 200–240 s	Coagulation factors ↓	FFP 20 ml/kg
R >14 min	ExTEM CT >100 s InTEM CT >240 s	Coagulation factors ↓↓	FFP 30 ml/kg
FF_{MA} 7–14 mm	FibTEM MCF 6–9 mm	Fibrinogen ↓	FFP 20 ml/kg or cryoprecipitate 3 ml/kg or fibrinogen concentrate 20 mg/kg
FF_{MA} 0–7 mm	FibTEM MCF 0–6 mm	Fibrinogen ↓↓	FFP 30 ml/kg or cryoprecipitate 5 ml/kg or fibrinogen concentrate 30 mg/kg
MA 45–49 mm and FF_{MA} >14 mm	ExTEM A_{10} 35–42 mm and FibTEM ≥10 mm ExTEM MCF <50 mm and FibTEM ≥10 mm	Platelets ↓	Platelets 5 ml/kg
MA <45 mm and FF_{MA} >14 mm	ExTEM A_{10} <35 mm and FibTEM ≥10 mm	Platelets ↓↓	Platelets 10 ml/kg
Ly30 >8%	ExTEM Li30 <94%	Hyperfibrinolysis	TXA 1–2 g IV or 10–20 mg/kg
Difference in R Hep TEG vs. standard TEG R >2 min	InTEM CT/HepTEM CT >1.25	Heparinazation	Protamine 50–100 mg or FFP 10–20 ml/kg

FFP, fresh frozen plasma; S, second; TXA, tranexamic acid.
Thrombelastograph (TEG): R, reaction time (normal range 3–8 min); MA, maximum amplitude (normal range 51–69 mm); Ly30, hyperfibrinolysis after 30 min (normal range <8%; FF_{MA}, functional fibrinogen MA (normal range 14–24 mm).
Thrombelastometry (ROTEM): CT, clotting time (normal range CT_{EXTEM} 38–79 s, CT_{INTEM} 100–240 s, CT_{HEPTEM} 100–240 s); A10, amplitude after 10 min (normal range $A10_{EXTEM}$ 43–65 mm, $A10_{FIBTEM}$ 7–23 mm); MCF, maximum clot firmness (normal range MCF_{EXTEM} 50–72 mm); Li30, hyperfibrinolysis after 30 min (normal range LI30 94–100%).

193. A 34-year-old man presents to the emergency department with severe abdominal pain and distension. On examination, his vital signs reveal a heart rate of 120 beats per minute and blood pressure of 80/50 mmHg. Abdominal examination reveals diffuse tenderness and distension. The plain abdominal radiograph is shown below. What is the most likely diagnosis?



A. Crohn's disease

B. Ulcerative colitis

C. Colorectal carcinoma

D. Toxic megacolon

Cerebellum
Get the balance right

194. A Female had a thorn prick 5 years ago. She presented with a slowly growing 2 × 2 cm verrucous lesion which on KOH mount showed the following image. Identify the condition?

A. Phaeohyphomycosis
B. Blastomycosis
C. Sporotrichosis
D. Chromoblastomycosis

1.



Cerebellum
Get the balance right

195. A 13-year-old girl is brought to the office for evaluation of acne. She began to develop acne over her forehead and chin 4 months ago. She used over-the-counter topical salicylic acid, but the acne did not improve. Now, the patient has painful acne over her face and upper chest. She has not had a menstrual period. The girl has grown 10.1 cm in the last year; her height and weight are at the 70th percentile for age. Vital signs are normal. Nodulocystic acne is present across the face and upper chest. There is no breast bud development. The abdomen is nontender and has no palpable masses. External pelvic examination shows the clitoris protruding from the clitoral hood and bilateral masses in the labia majora. Which of the following is the most likely diagnosis in this patient?

A. 5-alpha-reductase deficiency

B. Androgen insensitivity-syndrome

C. Nonclassical congenital adrenal hyperplasia

D. PCOD



196. A new antiarrhythmic drug is found to be effective against both atrial and ventricular arrhythmias. Its effect on action potential is shown in the Figure. The effect of this new drug is most similar to?

A. Lignocaine
B. Propanolol
C. Encainide
D. Quinidine



Ia

Cerebellum
Get the balance right

197. A 56-year-old man presented with the sudden onset of a severe headache. He was awake, but feeling uncomfortable because of headache. His initial plain CT head was normal. Lumbar puncture was performed and CSF was shown below. What is the next step?

A. Repeat plain CT head

B. Digital Subtraction Angiography

C. Ceftriaxone + Vancomycin

D. Aspirin



198. A 24-year-old man is brought to the emergency department due to weakness, lethargy, nausea, and dizziness. He took fifty 325-mg acetaminophen tablets approximately 6 hours ago. Vital signs are normal. Pupils are equal, round, and reactive to light; there is mild epigastric tenderness. Laboratory results include an elevated serum acetaminophen level. Intravenous infusion of N-acetylcysteine is started. This treatment is most likely to improve the patient's condition by which of the following mechanisms?

A. Activating hepatic glucuronidation enzymes

B. Competitively blocking drug receptor sites

C. Increasing the amount of intrahepatic glutathione

D. Increasing renal excretion of drug molecules

199. Identify the true statements about normal fetal development:

1. Endovascular type of extravillious trophoblasts invade the spiral arteries lumen
2. Blood in inferior vena cava has lower oxygen concentration compared to superior Vena cava
3. Functional closure of the fetal circulatory adjustments occurs in the following sequence: Ductus venosus > Foramen ovale > Ductus arteriosus
4. The main factor for the ductal closure postnatally is an increase in partial pressure of oxygen (paO2)

A. 1,2,3,4
B. 1,2,3
C. 1,3,4
D. 2,3,4

Cerebellum
Get the balance right

200. What does IP in the figure shown stands for?

A. Drug is an instant release preparation
B. Meant for intra pulmonary use
C. The standards according to Indian Pharmacopoeia
D. Drug tested preclinically via Intraperitoneal route

Cerebellum
Get the balance right

Thank You

Cerebellum

Get the balance right